

# PETITION ON HORMONAL CONTRACEPTIVES

William V. Williams, Joel Brind, Michael Manhart, Hanna Klaus, Angela Lanfranchi, Gerard Migeon,  
Michael Gaskins, Elvis Seman, Lester Ruppertsberger, Kathleen Raviele

CONTRACEPTIVE STUDY GROUP

## Contents

Preliminary Statement.....	4
Action Requested .....	4
Drugs which should be removed from the market: .....	4
Black box warnings that should be added to prescribing information .....	4
Additional safety information which should be added .....	5
List of Agents .....	6
Combined Estrogen-Progestin (EE-P) Pills.....	6
Combined EE-P Contraceptive Patch .....	10
Combined EE-P Contraceptive Ring .....	10
Progestin-Only Pills .....	11
Progestin-Only Injectable .....	11
Progestin-Only Implants.....	11
Progestin-Only IUS/IUD.....	11
Statement of Grounds .....	12
Risk of HIV Transmission .....	12
Epidemiological Evidence.....	12
Mechanistic Evidence.....	14
Summary and Conclusions: .....	14
Risk of HIV Transmission References .....	15
Cancer .....	17
Breast Cancer.....	17
Breast Cancer References .....	25
Cervical Cancer .....	30
Cervical Cancer References .....	33
Crohn’s Disease .....	34
Crohn’s Disease References .....	37
Ulcerative Colitis .....	39
Ulcerative Colitis References.....	42
Systemic Lupus Erythematosus .....	44
Systemic Lupus Erythematosus References.....	46
Risk of Depression, Mood Disorders, and Suicide .....	47
Depression, Mood Disorders, and Suicide References.....	51
Multiple Sclerosis.....	52
Multiple Sclerosis References.....	54

Interstitial Cystitis .....	55
Interstitial Cystitis References .....	55
Osteoporotic Bone Fractures .....	56
Osteoporotic Fracture References.....	61
Impact of Contraceptives on Body Mass .....	63
Impact of Contraceptives on Body Mass References .....	66
Urogenital Effects of Contraceptives.....	67
Urogenital Effects References .....	67
Venous Thromboembolism and Contraceptives.....	68
Venous Thromboembolism References.....	71
Atherosclerosis and Cardiovascular Events .....	74
Conclusion .....	77
Environmental Impact .....	77
Environmental Impact References .....	78
Economic Impact .....	80
HIV Costs .....	80
Breast Cancer.....	81
Cervical Cancer .....	82
Crohn’s Disease .....	83
Ulcerative Colitis.....	84
Systemic Lupus Erythematosus .....	85
Depression.....	85
Multiple Sclerosis.....	86
Interstitial Cystitis .....	87
Osteoporotic Bone Fracture Risk.....	87
Body Mass .....	88
Urogenital Effects .....	88
Venous Thromboembolism, Atherosclerosis and Cardiovascular Disease.....	88
Economic Costs References.....	89
Certification.....	93

## List of Tables

Table 1 – Meta-Analyses Evaluating Risk of HIV Transmission with Depot medroxyprogesterone acetate (DMPA) .....	12
Table 2 – Individual Studies of the Effects of DMPA HIV Transmission .....	13
Table 3 – Breast Cancer (Cohort Studies) .....	19
Table 4 – Breast Cancer (Case Control Studies) .....	21
Table 5 – Breast Cancer (Meta-Analyses) .....	24
Table 6 – Cervical Cancer .....	31
Table 7 – Individual Studies of the Effects of COCs on the Development of Crohn’s Disease .....	35
Table 8 – Individual Studies of the Effects of COCs on the Development of Ulcerative Colitis.....	40
Table 9 – Individual Studies of the Effects of COCs on the Development of Systemic Lupus Erythematosus..	45
Table 10 – Studies of Chemical Contraceptives and Depression, Mood Disorders and Suicides.....	48
Table 11 – Individual Studies of the Effects of COCs on the Development of Multiple Sclerosis .....	53
Table 12 – Individual Studies of the Effects of Contraceptives on the Development of Osteoporotic Fractures .....	57
Table 13 – Effect of Chemical Contraceptives on Weight Gain .....	64
Table 14 – Relative Risk of Venous Thromboembolism in Current Users of Different Combined Hormonal Contraceptives as Compared with Nonusers Unless Otherwise Specified .....	70
Table 15 – Relative Risk of Thrombotic Stroke and Myocardial Infarction among Users of Selected Types of Combined Oral Contraception with Ethinyl Estradiol at a Dose of 30 to 40 µg, as Compared with Nonusers, According to Duration of Use (from Lidegaard 2012). .....	75
Table 16 – Estimated Economic Impact of DMPA due to Increased Prevalence of HIV Infection .....	81
Table 17 – Estimated Economic Impact of COCs due to Increased Incidence of Breast Cancer .....	81
Table 18 – Estimated Economic Impact of COCs due to Increased Prevalence of Breast Cancer.....	82
Table 19 – Estimated Economic Impact of COCs due to Increased Incidence of Cervical Cancer.....	82
Table 20 – Estimated Economic Impact of COCs due to Increased Prevalence of Cervical Cancer.....	83
Table 21 – Estimated Economic Impact of COCs due to Increased Incidence of Crohn’s Disease .....	83
Table 22 – Estimated Economic Impact of COCs due to Increased Prevalence of Crohn’s Disease.....	83
Table 23 – Estimated Economic Impact of COCs due to Increased Incidence of Ulcerative Colitis .....	84
Table 24 – Estimated Economic Impact of COCs due to Increased Prevalence of Ulcerative Colitis .....	84
Table 25 – Estimated Economic Impact of COCs due to Increased Incidence of Systemic Lupus Erythematosus. ....	85
Table 26 – Estimated Economic Impact of COCs due to Increased Prevalence of Systemic Lupus Erythematosus. ....	85
Table 27 – Estimated Economic Impact of COCs due to Increased Prevalence of Depression .....	86
Table 28 – Estimated Economic Impact of POCs due to Increased Prevalence of Depression .....	86
Table 29 – Estimated Annual Economic Impact of COCs due to Increased Prevalence of Interstitial Cystitis..	87
Table 30 – Estimated Economic Impact of COCs due to Increased Annual Incidence of Vertebral Fractures..	87
Table 31 – Estimated Economic Impact of POCs due to Increased Annual Incidence of Vertebral Fractures..	88
Table 32 – Estimated Economic Impact of COCs due to Increased Incidence of Cardiovascular Disease .....	89
Table 33 – Cardiovascular Events in Women by Age Group.....	89
Table 34 – Cost of Cardiovascular Events in Women Attributable to COC use. ....	89

## Preliminary Statement

Hormonal contraceptives have been on the market for over 50 years and, while their formulations have changed, the basic mechanism of action has remained the same. During this time numerous studies have been performed documenting side effects, some of which appear over time, some within weeks or months, but all can have a serious impact on health. An effort was made to perform a series of comprehensive literature surveys to better understand immediate and long-term side effects of these agents. The results of this literature review have led to several recommendations. These recommendations are listed below with the documentation of the research noted on the following pages.

## Action Requested

### Drugs which should be removed from the market:

- Depot Medroxyprogesterone Acetate (DMPA)
  - Recommendation to remove from the market the injectable contraceptive Depot Medroxyprogesterone Acetate (DMPA; Depo Provera) based on conclusive evidence that it facilitates the transmission of HIV from men to women. Numerous alternatives are available.

### Black box warnings that should be added to prescribing information

- Breast Cancer
  - Combined estrogen-progestogen contraceptives (COCs, including oral, intravaginal and transdermal formulations) are acknowledged by IARC as Group I carcinogens. Substantial data supports an increased risk of breast cancer with the use of COCs. A black box warning should be added to the labeling of all COCs that they have been shown to increase the risk of breast cancer. Patient-related materials should also adequately convey this risk.
  - Progestogen-only contraceptives (POCs) have not been extensively studied, but one large registry study did show a significantly increased risk of breast cancer with use of POCs. Unless there is evidence to the contrary, a similar warning should be added to all POCs. Patient-related materials should also adequately convey this risk.
- Cervical Cancer
  - COCs have been linked to a significantly increased risk of cervical cancer. Similar data have been shown for POCs. A black box warning should be added to the labeling of all COCs and POCs that they have been shown to increase the risk of cervical cancer. Patient-related materials should also adequately convey this risk.
- Inflammatory Bowel Disease
  - Significantly higher risk for the development of inflammatory bowel disease, especially Crohn's disease, but also ulcerative colitis, has been shown for COCs. A black box warning should be added to the labeling of all COCs that their use is linked to a significantly increased risk for the development of inflammatory bowel disease. Patient-related materials should also adequately convey this risk.
- Systemic Lupus Erythematosus (SLE)
  - Significantly higher risk for the development of SLE has been shown for COCs in several studies, especially the best-designed, largest cohort studies. A black box warning should be added to the labeling of all COCs that their use is linked to a significantly increased risk of the development of SLE. Patient-related materials should also adequately convey this risk.
- Depression and Suicide

- Substantive evidence indicates there is a 25% risk of depression for women under 25 years of age especially within 6 months of starting COCs. A black box warning should be added to the labeling of all COCs that their use is linked to a significantly increased risk of the development of depression. Patient-related materials should also adequately convey this risk.
- The relative risk for suicide attempts ranges from 1.91 for COC's, to 2.29 for oral progestins, 2.58 for vaginal ring and 3.28 for patch among adolescents and young women – mean age 21 years – peaking within two months of onset of medication. A black box warning should be added to the labeling of all COCs that their use is linked to a significantly increased risk of suicide. Patient-related materials should also adequately convey this risk. Close monitoring is essential especially in the first year of use.
- Venous Thrombosis and Cardiovascular Events
  - The current black box warning regarding thrombotic events on some formulations, notes “WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS.” This is misleading and has shown to be misinterpreted by many women who infer that the increased risk only occurs with cigarette smoking and/or with being over 35 years of age. The warnings should be amended to state, “WARNING: INCREASED RISK OF SERIOUS CARDIOVASCULAR EVENTS INCLUDING BLOOD CLOTS.”
  - This warning should be required for hormonal birth control products including oral, intravaginal and transdermal formulations. The patient-related materials should clearly explain the genetic risk factors, other risk factors, and the signs and symptoms. This warning should be included in ALL direct-to-consumer advertising (television, print, radio, etc.).

#### Additional safety information which should be added

- Multiple Sclerosis (MS)
  - Significantly higher risk for the development of MS has been shown for COCs in several studies, especially the best-designed, largest case-control studies. A warning should be added to the labeling of all COCs that their use appears to be linked to a significantly increased risk of the development of MS. Patient-related materials should also adequately convey this risk.
- Bone Fractures
  - Use of POCs is clearly associated with a higher risk of bone fractures. A warning should be added to the labeling of all POCs that their use is linked to a significantly increased risk of the development of bone fractures. Patient-related materials should also adequately convey this risk.
  - Protracted use of COCs has been associated with an increased risk of bone fractures. A warning should be added to the labeling of all COCs that their prolonged use may be linked to a significantly increased risk of the development of bone fractures. Patient-related materials should also adequately convey this risk.
- Body Mass Effects
  - For ANY progestin-releasing IUD:
    - Add to professional label in side effects/precautions:
      - Progestin-releasing IUDs (IUCs) have demonstrated in clinical trials to significantly increase % fat body mass with a corresponding decrease in % lean body mass over 1 year of use.
    - Add to patient-related materials:
      - Use of (Brand name) may increase the percent of fat in your body while decreasing the percent of lean body mass; this change in body composition is

known to increase risk of other serious conditions such as diabetes and cardiovascular problems.

- This warning should be included in all direct-to-consumer advertising (television, print, radio, etc.) as it demonstrates use of IUCs may contribute to other serious chronic health conditions.
- Similar labeling should be considered for progestin-only contraceptives. Although the current evidence is less, it tends in the same direction.
- Urogenital Problems
  - Interstitial Cystitis: Significantly higher risk for the development of interstitial cystitis has been shown for COCs in two studies. A warning should be added to the labeling of all COCs that their use appears to be linked to a significantly increased risk of the development of interstitial cystitis. Patient-related materials should also adequately convey this risk.
  - COCs have also been linked to an increased risk of bacteriuria, urinary tract infections, bladder trabeculation, vulvovaginal candidiasis, vaginal dryness, vulvar vestibulitis, and Female Sexual Dysfunction (FSD) caused by OC-induced dyspareunia and reduced sexual desire and libido. These risks should be adequately conveyed in the prescribing information, especially FSD where there is substantial literature evidence.

### List of Agents

A list of the agents discussed is shown below. Other than Depot Medroxyprogesterone Acetate (DMPA; Depo Provera) we refer in general to COCs (which refers to all combined estrogen-progestogen contraceptive formulations) and POCs (which refers to all progestin-only contraceptive formulations) regardless of the route of administration (e.g. oral, intravaginal, transdermal, implants, IUS/IUD, etc.).

#### Combined Estrogen-Progestin (EE-P) Pills

OVCON-35

FEMCON 35

FEMCON FE

BALZIVA 28

BRIELLYN 28

PHILITH

GILDAGIA

VYFEMLA

NEXESTA FE

and generic therapeutic equivalents

BREVICON

MODICON 28

NORMINEST FE

NORTREL 0.5/35-28

WERA

CYCLAFEM

CYONANZ

and generic therapeutic equivalents

GENERESS

KAITLIB FE

and generic therapeutic equivalents

NORINYL 1+35 28-DAY TABLETS  
ORTHO-NOVUM 1/35 28 TABLETS  
ALYACEN 1/35  
ARANELLE  
CYCLAFEM 1/35  
DASETTA 1/35  
NORTREL 1/35-28  
NYLIA 1/35  
PIRMELLA 1/35

and generic therapeutic equivalents

ORTHO-NOVUM 7/7/7-28  
ALYACEN 7/7/7  
CYCLAFEM 7/7/7  
DASETTA 7/7/7  
NORTREL 7/7/7  
NYLIA 7/7/7  
PIRMELLA 7/7/7

TRI-NORINYL 28-DAY  
ARANELLE

NORINYL 1+50 28-DAY

LOESTRIN 21 1/20  
LOESTRIN 21 1/20 FE  
MINASTRIN 24 FE  
TAYTULLA  
MIBELAS 24 FE

MICROGESTIN 1/20  
MICROGESTIN FE 1/20  
JUNEL 1/20  
GILDESS 1/20 and GILDESS FE 1/20  
LARIN 1/20 and LARIN FE 1/20  
BLISOVI 1/20 and BLISOVI FE 1/20  
AUROVELA 1/20 and AUROVELA 1/20 FE  
HAILEY 1/20 and HAILEY FE 1/20  
and generic therapeutic equivalents

LOESTRIN 21 1.5/30  
LOESTRIN FE  
MICROGESTIN 1.5/30  
MICROGESTIN FE  
AUROVELA 1.5/30



AUROVELA FE 1.5/30  
BLISOVI FE 1.5/30  
GILDESS 1.5/30  
GILDESS FE 1.5/30  
JUNEL 1.5/30  
JUNEL FE  
LARIN 1.5/30  
LARIN FE

ESTROSTEP 21  
ESTROSTEP FE  
TRI-LEGEST 21  
TRI-LEGEST FE  
and generic therapeutic equivalents

ZOVIA 1/35E-28  
KELNOR  
and generic therapeutic equivalents

LOW-OGESTREL-28  
CRYSSELLE  
ELINEST

OGESTREL 0.5/50-28

LoSEASONIQUE  
LO SIMPESSÉ  
and generic therapeutic equivalents

ALESSE  
LEVLITE  
LESSINA-28  
AVIANE-28  
BALCOLTRA  
AFIRMELLE  
FALMINA  
ORSYTHIA  
VIENVA  
and generic therapeutic equivalents

QUARTETTE—91-DAY  
FAYOSIM

SEASONALE  
INTROVALE  
ALTAVERA  
AYUNA

QUASENSE  
SETLAKIN  
LEVORA 0.15/30-28  
KURVELO  
PORTIA-28  
MARLISSA

SEASONIQUE  
ASHLYNA  
DAYSEE  
JAIMIESS  
SIMPESSE  
and generic therapeutic equivalents

TRIVORA-28  
ENPRESSE-28  
LEVONEST  
ELIFEMME  
MYZILRA  
and generic therapeutic equivalents

DESOGEN  
EMOQUETTE  
ENSKYCE  
ISIBLOOM  
KALLIGA  
and generic therapeutic equivalents

KARIVA  
KIMIDESS  
VIORELE  
PIMTREA  
VOLNEA  
BEKYEE  
and generic therapeutic equivalents

CYCLESSA  
VELIVET  
and generic therapeutic equivalents

ORTHO-CYCLEN-28  
SPRINTEC  
PREVIFEM  
MONO-LINYAH  
ESTARYLLA  
MILI  
and generic therapeutic equivalents

ORTHO TRICYCLEN 28  
TRI-SPRINTEC  
TRIPREVIFEM-28  
TRI-LINYAH  
TRI-ESTARYLLA  
TRI-MILI  
and generic therapeutic equivalents

ORTHO TRI-CYCLEN LO  
TRI-PREVIFEM  
TRI LO SPRINTEC  
TRI-LO-ESTARYLLA  
TRI-LO-MILI  
and generic therapeutic equivalents

YAZ  
LORYNA  
NIKKI  
MELAMISA  
LO-ZUMANDIMINE  
and generic therapeutic equivalents

BEYAZ  
and generic therapeutic equivalents

YASMIN 28  
SYEDA  
YAELA  
ZUMANDIMINE  
and generic therapeutic equivalents

SAFYRAL

NATAZIA

Combined EE-P Contraceptive Patch  
ORTHO EVRA  
XULANE

Combined EE-P Contraceptive Ring  
NUVARING

Progestin-Only Pills

MICRONOR TABLETS

NOR-QD TABLETS

CAMILA

ERRIN

HEATHER

JENCYCLA

INCASSIA

and generic therapeutic equivalents

Progestin-Only Injectable

DEPO PROVERA

Progestin-Only Implants

JADELLE

NEXPLANON

Progestin-Only IUS/IUD

MIRENA IUS

LILETTA IUD

SKYLA IUD

KYLEENA IUD

## Statement of Grounds

### Risk of HIV Transmission

One of the most common forms of steroidal contraception is the injectable contraceptive: Depot medroxyprogesterone acetate (DMPA). DMPA is highly effective and requires only quarterly injections, as opposed to daily oral ingestion. As a long-acting type of effective contraceptive, it is not unique, as there are other injectable or implantable contraceptives in wide use, e.g., norethisterone enanthate (NET), as well as other delivery systems such as vaginal rings and patches.

However, evidence began emerging in the 1990s, which has become compelling in recent years, that DMPA is unique among contraceptives in its property of facilitating the transmission of HIV. This dangerous characteristic has been abundantly and unequivocally documented through several lines of evidence which are summarized below:

### Epidemiological Evidence

- A. Four meta-analyses (3 reports) were published in 2015. Each used different inclusion criteria and compiled the data on different numbers of studies, yet all 4 came up with essentially the same result of significantly increased risk of male-to-female HIV transmission in women using DMPA (Table 1).

*Table 1 – Meta-Analyses Evaluating Risk of HIV Transmission with Depot medroxyprogesterone acetate (DMPA)*

Meta-analysis	# Included studies	Pooled Adj. OR or HR (95% CI)
Ralph et al. 2015	10 (longitudinal)	HR 1.40 (1.16–1.69)
Morrison et al. 2015	18 (longitudinal)	HR 1.50 (1.24–1.83)
Brind et al. 2015	8 (cross-sectional)	OR 1.41 (1.15–1.73)
Brind et al. 2015	16 (longitudinal)	HR 1.49 (1.28–1.73)

- B. Ten primary studies (all longitudinal, published between 2003 and 2014, listed in Table 2 below) were methodologically robust enough to meet the inclusion criteria of all 3 published reviews.

Table 2 – Individual Studies of the Effects of DMPA HIV Transmission

Study	Yr.(s) of study	Pop. size	Nation and locale	Subject source	Months of follow-up	Follow-up interval (months)	Type of data shown	HR or IRR (95% CI)	Weight (%)
Crook 2014	2005–2009	8,663	S Africa, Uganda, Tanzania, Zambia	Microbicide trial sero-disc. couples	12	1	Inv. Prob. W'ted HR	1.45 (1.09–1.93)	16.39
McCoy 2013	2003–2007	4,913	South Africa, Zimbabwe	Diaphragm/gel HIV prev. trial	24	3	MV HR	1.22 (0.85–1.76)	13.20
Morrison 2012	2004–2007	5,567	South Africa	General population	9–24	3	MSM HR	1.27 (0.93–1.73)	15.32
Wand 2012	Not reported	2,236	Durban, S. Africa	>90% from microbicide trial	Not reported	3	MV HR	2.02 (1.37–2.99)	12.22
Heffron 2012	2004–2010	3,790	7 African nations	Sero-discordant couples	12–24	3	MSM HR	3.93 (1.38–11.21)	2.81
Morrison 2007	1999–2004	6,109	Uganda, Zimbabwe, Thailand	Family planning clinics	21.5	3	MSM HR	1.25 (0.88–1.77)	13.86
Myer 2007	2000–2004	4,073	Cape Town, So. Africa	General population	24	6,6, & 12	MV IRR	0.75 (0.33–1.69)	4.36
Kleinschmidt 2007	1999–2001	551	Orange Farm, So. Africa	Family planning clinic	12	3	MV HR	0.46 (0.06–3.66)	0.78
Baeten 2007	1993–1997	779	Mombasa, Kenya	CSW	120	1	MV HR	1.73 (1.28–2.34)	15.69
Kiddugavu 2003	1994–1999	5,117	Rakai, Uganda	General population	31	10	IRR, MLR	0.84 (0.41–1.72)	5.37

Importantly, no consistent association has emerged with regard to oral contraceptives or other injectable or implantable contraceptives and the facilitation of HIV transmission.

#### Mechanistic Evidence

- A. *In vivo* evidence of increased HIV transmission: Heffron et al. (2012) reported the increased presence of HIV-1 RNA in genital fluids of women using DMPA.
- B. *In vitro* evidence of increased HIV replication at the cellular level: Maritz et al. (2018) reported experimental evidence of increased replication of HIV in human blood monocytes with medroxyprogesterone acetate (MPA).
- C. Experimental evidence of agonistic binding to the glucocorticoid receptor (GR) as the mechanism for DMPA's immunosuppression: over the last 15 years, abundant experimental evidence of cytotoxic and immunosuppressive action of DMPA via its agonistic binding to the GR of human leukocytes has been reported (Schindler 2003; Hapgood and Tomasicchio 2010, Hapgood 2014.) Thus, Huijbregts et al. (2014) reported experimental evidence of immunosuppression of human T cells in vitro by MPA. Tomasicchio et al. (2013) reported experimental evidence of increased human T-cell destruction in vitro via the glucocorticoid receptor (GR) with MPA. Hapgood et al. (2014) reported:  
*“that MPA, unlike NET and progesterone, represses inflammatory genes in human PBMCs (peripheral blood mononuclear cells) in a dose-dependent manner, via the glucocorticoid receptor (GR), at concentrations within the physiologically relevant range. These and published results collectively suggest that the differential GR activity of MPA versus NET may be a mechanism whereby MPA, unlike NET or progesterone, differentially modulates HIV-1 acquisition and pathogenesis in target cells where the GR is the predominant steroid receptor expressed.”*
- D. Evidence of mechanism of MPA action at the gene expression level: experimental evidence of MPA-mediated suppression of inflammatory genes via GR in cultured human cells (Govender 2014) demonstrated the suppression of inflammatory genes in cultured human endocervical cells.

#### Summary and Conclusions:

DMPA, in contrast to all other steroidal contraceptives, has now conclusively been demonstrated to significantly increase the risk of HIV transmission from infected men to women. The robust epidemiological association has been supported by *in vivo* evidence of increased HIV RNA in the female genital tracts of women using DMPA. Moreover, abundant experimental evidence has shown that MPA, due to its agonistic binding of the GR, specifically represses the innate immune responses of both circulating human leukocytes and endocervical cells and allows for increasing HIV replication. The demonstration in the literature of the chain of causation is therefore compelling.

In the United States, where the availability of a wide range of contraceptive drugs and devices is virtually universal, and where, among these contraceptive choices, one and only one particular method—DMPA—is now known to increase the transmission of an often-fatal viral infection (HIV/AIDS), there can be no justification for such a drug's continued availability in the marketplace. It should be removed from the marketplace by the FDA without further delay.

## Risk of HIV Transmission References

Baeten JM, Benki S, Chohan V, Lavreys L, McClelland RS, Mandaliya K, Ndinya-Achola JO, Jaoko W, and Overbaugh J. Hormonal contraceptive use, herpes simplex virus infection, and risk of HIV-1 acquisition among Kenyan women. *AIDS* 2007; 21:1771–1777.

Brind J, Condly SJ, Mosher SW, Morse AR, and Kimball J. Risk of HIV infection in Depot Medroxyprogesterone Acetate (DMPA) users: a systematic review and meta-analysis. *Issues Law Med* 2015; 30:129–139.

Crook AM, Ford D, Gafos M, Hayes R, Kamali A, Kapiga S, Nunn A, Chisembele M, Ramjee G, Rees H, and McCormack S. Injectable and oral contraceptives and risk of HIV acquisition in women: an analysis of data from the MDP301 trial. *Hum Reprod* 2014; 29:1810–1817.

Govender Y, Avenant C, Verhoog NJ, Ray RM, Grantham NJ, Africander D, and Hapgood JP. The injectable-only contraceptive medroxyprogesterone acetate, unlike norethisterone acetate and progesterone, regulates inflammatory genes in endocervical cells via the glucocorticoid receptor. *PLoS One* 2014; 9:e96497.

Hapgood JP and Tomasicchio M. Modulation of HIV-1 virulence via the host glucocorticoid receptor: towards further understanding the molecular mechanisms of HIV-1 pathogenesis. *Arch Virol* 2010; 155:1009-1019. Doi: 10.1007/s00705-010-0678-0.

Hapgood JP, Ray RM, Govender Y, Avenant C, and Tomasicchio M. Differential glucocorticoid receptor-mediated effects on immunomodulatory gene expression by progestin contraceptives: implications for HIV-1 pathogenesis. *Am J Reprod Immunol* 2014; 71(6):505–512. Doi: 10.1111/aji.12214.

Heffron R, Donnell D, Rees H, Celum C, Mugo N, Were E, de Bruyn G, Nakku-Joloba E, Ngure K, Kiarie J, Coombs RW, Baeten JM; Partners in Prevention HSV/HIV Transmission Study Team. Use of hormonal contraceptives and risk of HIV-1 transmission: A prospective cohort study. *Lancet Infect Dis* 2012; 12:19–26.

Huijbregts RP, Michel KG and Hel Z. Effect of progestins on immunity: medroxyprogesterone but not norethisterone or levonorgestrel suppresses the function of T cells and pDCs. *Contraception* 2014 Aug; 90(2):123–129. Doi: 10.1016/j.contraception.2014.02.006.

Kiddugavu M, Makumbi F, Wawer MJ, Serwadda D, Sewankambo NK, Wabwire-Mangen F, Lutalo T, Meehan M, Xianbin, Gray RH; Rakai Project Study Group. Hormonal contraceptive use and HIV-1 infection in a population-based cohort in Rakai, Uganda. *AIDS* 2003; 17:233–240.

Kleinschmidt I, Rees H, Delany S, Smith D, Dinat N, Nkala B, and McIntyre JA. Injectable progestin contraceptive use and risk of HIV infection in a South African family planning cohort. *Contraception* 2007; 75:461–467.

Maritz MF, Ray RM, Bick AJ, Tomasicchio M, Woodland JG, Govender Y, Avenant C, and Hapgood JP. Medroxyprogesterone acetate, unlike norethisterone, increases HIV-1 replication in human peripheral blood mononuclear cells and an indicator cell line, via mechanisms involving the glucocorticoid receptor, increased CD4/CD8 ratios and CCR5 levels. *PLoS One* 2018 <https://doi.org/10.1371/journal.pone.0196043>.



McCoy SI, Zheng W, Montgomery ET, Blanchard K, van der Straten A, de Bruyn G, and Padian NS. Oral and injectable contraception use and risk of HIV acquisition among women in sub-Saharan Africa. *AIDS* 2013; 27:1001–1009.

Morrison CS, Richardson BA, Mmiro F, Chipato T, Celentano DD, Luoto J, Mugerwa R, Padian N, Ruggao S, Brown JM, Cornelisse P, Salata RA; Hormonal Contraception and the Risk of HIV Acquisition (HC-HIV) Study Group. Hormonal contraception and the risk of HIV acquisition. *AIDS* 2007; 21:85–95.

Morrison CS, Skoler-Karpoff S, Kwok C, Chen PL, van de Wijgert J, Gehret-Plagianos M, Patel S, Ahmed K, Ramjee G, Friedland B, and Lahteenmaki P.. Hormonal contraception and the risk of HIV acquisition among women in South Africa. *AIDS* 2012; 26:497–504.

Morrison CS, Chen P-L, Kwok C, Baeten JM, Brown J, Crook AM, Van Damme L, Delany-Moretlwe S, Francis SC, Friedland BA, Hayes RJ, Heffron R, Kapiga S, Karim QA, Karpoff S, Kaul R, McClelland RS, McCormack S, McGrath N, Myer L, Rees H, van der Straten A, Watson-Jones D, van de Wijgert JH, Stalter R, and Low N. Hormonal contraception and the risk of HIV acquisition: an individual participant data meta-analysis. *PLoS Med* 2015; 12:e1001778. doi: 10.1371/journal.pmed.1001778.

Myer L, Denny L, Wright TC, and Kuhn L. Prospective study of hormonal contraception and women's risk of HIV infection in South Africa. *Int J Epidemiol* 2007; 36:166–174.

Ralph LJ, McCoy SI, Shiu K and Padian NS. Hormonal contraceptive use and women's risk of HIV acquisition: a meta-analysis of observational studies. *Lancet Infect Dis* 2015; 15:181–189. Doi: 10.1016/S1473-3099(14)71052–1057.

Schindler AE, Campagnoli C, Druckmann R, Huber J, Pasqualini JR, Schweppe KW, and Thijssen JH. Classification and pharmacology of progestins. *Maturitas* 2003; 46 (Suppl 1):S7-S16. Doi: 10.1016/j.maturitas.2003.09.014.

Tomasicchio, Michele, Avenant C, Du Toit A, Ray RM, and Hapgood JP. The Progestin-Only Contraceptive Medroxyprogesterone Acetate, but Not Norethisterone Acetate, Enhances HIV-1 Vpr-Mediated Apoptosis in Human CD4+ T Cells through the Glucocorticoid Receptor. *PLoS One* 2013; 8:e62895.

Wand H and Ramjee G. The effects of injectable hormonal contraceptives on HIV seroconversion and on sexually transmitted infections. *AIDS* 2012; 26:375–380.

## Cancer

Papers were accessed from a PubMed literature review as noted (Williams 2018). Each paper was rated based on the parameters noted in the STROBE statement (von Elm et al. 2007).

### Breast Cancer

Breast cancer is the most commonly diagnosed cancer (excluding non-melanoma skin cancers) in women in developed nations, including the U.S., with 1.7 million cases diagnosed worldwide annually. It accounts for 20% of all cancers in women.<sup>1</sup> According to the Surveillance, Epidemiology and End Results (SEER) statistics<sup>2</sup>, it is estimated that there are about 3,418,000 women with invasive breast cancer in the USA as well as over 60,000 cases of in situ cancers. There will be about 266,000 new cases of breast cancer in 2018, accounting for 15.3% of all new cancer cases, with about 41,000 deaths, accounting for 6.7% of all cancer deaths. Nulliparity or late childbearing and high body mass index are risk factors for breast cancer as is exposure to COCs and HRT. Any risk factors that are controllable should be minimized. The data for breast cancer is shown split into cohort studies (Table 3), case control studies (Table 4) and meta-analyses (Table 5).

The carcinogenicity of combined estrogen-progestogen contraceptives was evaluated by IARC working groups initially in 1998 (monograph published in 1999) and again in 2005 (monograph published in 2007). This was most recently updated with studies published through May 2008 (IARC 2012). Since that time, several important studies have been published, most of which are supportive of the IARC classification of COCs as Group I carcinogens and in agreement with the IARC evaluation of specific cancer types. In addition, several important studies have been published evaluating COCs and their cancer risk. In 2002 the National Toxicology Program added steroidal estrogen as a known human carcinogen (Report on Carcinogens, Fourteenth Edition available at <https://ntp.niehs.nih.gov/pubhealth/roc/index-1.html>).

In agreement with IARC the recent data confirms an increased risk of breast cancer with use of COCs (Table 2). After 2005, there continue to be studies demonstrating the significant risk of breast cancer with hormonal contraception. In January 2006, the *New England Journal of Medicine* published a review article which found estrogen-progestin drugs increased breast cancer risk (Yager 2006). In October 2006 the *Mayo Clinic Proceedings* published a meta-analysis confirming estrogen-progestin drugs increase premenopausal breast cancer (Kahlenborn 2006).

The studies that looked at recent use (within 1–5 years) or current use of COCs in premenopausal women showed the most dramatic increased risk for breast cancer. In a case control study, women ages 20-49 years with use of COCs within a year had an increased risk of breast cancer (OR, 1.5; 95% CI, 1.3–1.9) (Beaber 2014). The same study showed an increase in risk depending on the formulation with triphasic COCs carrying a markedly increased risk (OR, 3.1; 95% CI, 1.4–4.7). In another large case control study of women ages 20-45 years, use of COCs for a year or more resulted in a 2.5-fold increased risk of triple-negative breast cancer (95% CI 1.4–4.3) but not for the receptor-positive breast cancers. In the same study, women 40 years or younger with a year or more use of COCs had a higher relative risk of triple-negative breast cancer (RR, 4.2; 95% CI, 1.9–9.3) (Dolle 2009). A cohort study of over 35,000 postmenopausal women found a significantly increased risk of breast cancer in women on hormone replacement therapy (HRT) if they had used COCs in the past (RR, 2.45; 95% CI, 1.92–3.12) as compared with never users (RR, 1.67; 95% CI, 1.32–2.12) (Lund 2007). There also appears to be an increased risk for African American women on COCs within the past five years for ER+ cancers (OR, 1.46, 95% CI, 1.18–1.81), for ER- cancers (OR, 1.57; 95% CI, 1.22–1.43) and for triple-negative

<sup>1</sup> <https://www.wcrf.org/int/cancer-facts-figures/data...cancers/breast-cancer-statistics>.

<sup>2</sup> <https://seer.cancer.gov/statfacts/html/breast.html>.

cancers (OR, 1.78; 95% CI, 1.25–2.53) with the risk of ER+ cancers continuing for 15–19 years after stopping the COCs (Bethea 2015).

In a French study (DeLort 2007) of 934 women who developed breast cancer, the use of COCs increased the risk of early development of breast cancer (OR, 1.84; 95% CI, 1.38–2.44). However, initiating COCs after age 23 reduced the risk (OR, 0.52; 95% CI, 0.34–0.79). Use of the levonorgestrel-releasing IUD, commonly used to treat abnormal bleeding in the perimenopause, increased the risk of developing breast cancer in postmenopausal women (OR, 1.48, 95% CI, 1.10–1.99) (Heikkinen 2016). The risk varies with the formulation as current use of a triphasic pill containing levonorgestrel carries an excess risk of causing breast cancer (RR, 3.05; 95% CI, 2.00–4.66) (Hunter 2010). In a large prospective cohort study of 1.8 million Danish women ages 15 to 49, enrolled and followed from 1995 to 2012 through various national registries, the risk of breast cancer among current or recent users increased depending on length of use from RR, 1.09 with less than one year of use (95% CI, 0.96–1.23) to an RR, 1.38 (95% CI, 1.26–1.51) for more than 10 years of use (Mørch 2017). They found the increased risk persisted after discontinuing use if COCs were used for 5 years or more. These investigators also found an increased risk in current or recent use of the progestogen-only intrauterine device (RR, 1.21; 95% CI, 1.11–1.33).

In most Western countries, 5% to 10% of all breast cancer cases are due to a main genetic cause: mutations of the BRCA1 and BRCA2 genes constitute 90% of hereditary breast cancer cases (Mehrgou 2016). These women are often begun on COCs at an early age to reduce their risk of ovarian cancer. However, in a case control study of 2,492 matched pairs of women with the *BRCA1* gene, COC use was associated with an increased risk of early onset breast cancer if begun under the age of 20 (OR, 1.45; 95% CI, 1.20–1.75) (Kotsopoulos 2014) and the risk increased by 11% for each additional year of use.

More recent publications include data from some very recent, large cohort studies (Mørch 2017, Heikkinen 2016, Poosari 2014) with RRs ranging from 1.2 to 1.37. Since breast cancer is by far the most common cancer in women, affecting 1 in 8 women at some time during their lives, this translates into a substantial number of additional cancer cases. In addition, a large registry study of POCs (Soini 2014) also showed an increased RR for breast cancer of 1.19. Increased duration of use also increases the risk of breast cancer for COCs as does use early in life (Mørch 2017).

Table 3 – Breast Cancer (Cohort Studies)

Study	Study Design	OR <sup>1</sup> Ever Use	RR <sup>2</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Mørch et al. 2017	Cohort		1.2 <sup>3</sup> (1.14–1.26)					1,797,932	* <sup>4</sup>	100%
Heikkinen et al. 2016	Cohort		1.37 (1.12–1.68)					7,000	20,000	100%
Lund et al. 2007	Cohort		1.33 (1.11–1.59)					11,777	23,676	96%
Poosari et al. 2014	Cohort		1.31 (0.65–2.65)					70	11,344	92%
Phipps et al. 2011	* <sup>5</sup>		0.80 <sup>6</sup> (0.68–0.94)					5,194		92%
Brohet et al. 2007 <sup>7</sup>	Cohort		1.47 (1.16–1.87)					846	747	88%
Thorbjarnardottir et al. 2014	Cohort		1.32 (1.02–1.70)					654	16,928	84%
Samson et al. 2017	Cohort		1.80 <sup>8</sup> (1.29–2.55)					4816		83%
Rosenberg et al. 2010	Cohort		1.65 (1.19–2.30)					789	53,848	83%
Silvera et al. 2005	Cohort		0.88 <sup>9</sup> (0.73–1.07)					1,707	25,611	78%
Hunter et al. 2010	Cohort		1.12 (0.95–1.33)		1.33 (1.03–1.73)			1,344	115,264	73%
			1.42 <sup>10</sup> (1.05–1.94)							

<sup>1</sup> OR = odds ratio (95 % confidence interval).

<sup>2</sup> RR = relative risk (95 % confidence interval).

<sup>3</sup> Initiation before age 20, greater than 10 years of use and evaluation within 5 yrs. of stopping further increased the risk.

<sup>4</sup> Entire population of Denmark was the cohort.

<sup>5</sup> Concurrent randomized clinical trials and an observational study.

<sup>6</sup> Hazard ratio shown. Note that women started COCs after age 25, had been off COCs for many years.

<sup>7</sup> Evaluation in patients carrying BRCA mutations. Hazard ratios shown.

<sup>8</sup> Hazard ratio shown.

<sup>9</sup> Hazard ratio shown.

<sup>10</sup> Eight or more years of use.

Study	Study Design	OR <sup>1</sup> Ever Use	RR <sup>2</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
			3.05 <sup>11</sup> (2.00–4.66)							
Trivers et al. 2007 <sup>12</sup>	Cohort			1.57 (0.95–2.61)				292 <sup>13</sup>	1,264 <sup>14</sup>	67%

---

<sup>11</sup> Levonorgestrel containing combined oral contraceptives.

<sup>12</sup> Looked at mortality in patients with breast cancer over 8-10 years depending on whether they were on COCs at the time of diagnosis or within one year.

<sup>13</sup> Deaths.

<sup>14</sup> Total cohort.

Table 4 – Breast Cancer (Case Control Studies)

Study	Study Design	OR <sup>15</sup> Ever Use	RR <sup>16</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Dolle et al. 2009	Case control	2.5 (0.9-5.24)		4.2 (1.9-9.3)				898	961	100%
Lee et al. 2008	Case Case <sup>17</sup>	0.68 (0.33-1.38)						94	444	100%
Sweeney et al. 2007	Case control	1.27 (0.99-1.63)						2,318	2,515	100%
Beaber et al. 2014b	Case control	1.5 (1.1-2.2)						985	882	100%
Li et al. 2012 <sup>18</sup>	Case control	2.2 (1.2-4.2)						1,028	919	96%
Beaber et al. 2014a	Case control			1.5 <sup>19</sup> (1.3-1.9)				1,102	21,952	96%
Ichida et al. 2015	Case control			0.45 (0.22-0.90)				155	12,333	96%
Ma et al. 2010	Case control	2.87 <sup>20</sup> (1.44-5.74)						1,197	2,015	95%
Folger et al. 2007	Case control	1.0 <sup>21</sup> (0.8-1.1)						4575	4682	92%
Jernstrom et al. 2005	Case control					2.10 (1.32-3.33)		245	745	92%
Kotsopoulos et al. 2014 <sup>22</sup>	Case control	1.45 <sup>23</sup> (1.20-1.75)						2,492	2,492	88%
		1.19 <sup>24</sup> (0.99-1.42)								
Figueiredo et al. 2010 <sup>25</sup>	Case control					2.38 (0.72-7.83)		705	1,398	86%

<sup>15</sup> OR = odds ratio (95 % confidence interval).

<sup>16</sup> RR = relative risk (95 % confidence interval).

<sup>17</sup> BRCA1 and BRCA2 carriers with breast cancer.

<sup>18</sup> Population-based case-control of women 20-44 yo with recent DMPA use for at least 12 months.

<sup>19</sup> Use within the past year of COCs increases risk of breast cancer.

<sup>20</sup> Triple negative breast cancer if less than 18 yo on COCs.

<sup>21</sup> Evaluated short-term use only.

<sup>22</sup> Study of BRCA+ patients.

<sup>23</sup> <20 years old.

<sup>24</sup> 20-25 years old.

<sup>25</sup> Evaluation of BRCA1 and BRCA2 carriers; controls with unilateral breast cancer compared with contralateral cases.

Study	Study Design	OR <sup>15</sup> Ever Use	RR <sup>16</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Veneroso et al. 2008	Case Case <sup>26</sup>	1.12 (1.03-1.23)						116	99	86%

Study	Study Design	OR <sup>27</sup> Ever Use	RR <sup>28</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Ma et al. 2006	Case control	1.27 <sup>29</sup> (0.75-2.14)				0.76 (0.49-1.18)		1,366	440	84%
		0.76 <sup>30</sup> (0.49-1.18)								
Rosenberg et al. 2008	Case control	1.5 <sup>31</sup> (1.2-1.8)						907	1,711	83%
Haile et al. 2006	Case control	0.77 <sup>32</sup> (0.53-1.12)						195	497	83%
		1.62 <sup>33</sup> (0.90-2.92)						128	307	
Milne et al. 2005	Case control	1.52 (1.22-1.91)						1156	815	83%
Amadou et al. 2013	Case control	1.68 (0.67-4.21)						1,000	1,074	75%
Ozmen et al. 2009	Case control	0.60 (0.48-0.74)						1,492	2,167	74%
Delort et al. 2007	Population based <sup>34</sup>	1.84 <sup>35</sup> (1.38-2.44)						934		71%
Beji et al. 2006	Case control	1.98 (1.38-2.85)						405	1,050	63%
Veisy et al. 2015	Case control	2.11 (1.44-3.08)						235	235	63%
Adams-Campbell et al. 2010	Case control	2.83 (1.87-4.24)						321	321	58%

<sup>26</sup> Comparison of more aggressive with less aggressive cases.

<sup>27</sup> OR = odds ratio (95 % confidence interval).

<sup>28</sup> RR = relative risk (95 % confidence interval).

<sup>29</sup> ER-/PR-

<sup>30</sup> ER+/PR+

<sup>31</sup> OR for 5+ years of use.

<sup>32</sup> BRCA1+ patients.

<sup>33</sup> BRCA2+ patients.

<sup>34</sup> Population-based study of early onset breast cancer.

<sup>35</sup> OR for developing breast cancer 2 years earlier than non-users.

Study	Study Design	OR <sup>27</sup> Ever Use	RR <sup>28</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Lumachi et al. 2010	Retrospective Review	2.06 (1.14-3.70)						404	408	33%



Table 5 – Breast Cancer (Meta-Analyses)

Study	Study Design	OR <sup>36</sup> Ever Use	RR <sup>37</sup> Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Kahlenborn et al. 2006 <sup>38</sup>	Meta-analysis	1.19 (1.09-1.29)						18,406	27,677	91%
		1.29 <sup>39</sup> (1.20-1.40)								
		1.24 <sup>40</sup> (0.92-1.67)								
		1.44 <sup>41</sup> (1.28-1.62)								
Bethea et al. 2015	Meta-analysis	1.46 <sup>42</sup> (1.18-1.81)						1,848	10,044	85%
		1.57 <sup>43</sup> (1.22-1.43)						1,043	10,044	
		1.78 <sup>44</sup> (1.25-2.53)						494	10,044	
Zhu et al. 2012	Meta-analysis	1.08 <sup>45</sup> (0.99-1.17)								54%
Friebel et al. 2014 <sup>46</sup>	Meta-analysis	1.36 <sup>47</sup> (0.99-1.88)								27%
		1.51 <sup>48</sup> (1.10-2.08)								
Moorman et al. 2013	Meta-analysis	1.21 <sup>49</sup> (0.93-1.58)								

<sup>36</sup> OR = odds ratio (95 % confidence interval).

<sup>37</sup> RR = relative risk (95 % confidence interval).

<sup>38</sup> Limited to case-control studies from 1980-2004.

<sup>39</sup> Parous women.

<sup>40</sup> Nulliparous women.

<sup>41</sup> Use before first full term pregnancy among parous women.

<sup>42</sup> ER+

<sup>43</sup> ER-

<sup>44</sup> Triple negative.

<sup>45</sup> For each 5 years on COCs the risk increased by 7%, but statistical significance not achieved.

<sup>46</sup> Study limited to BRCA1 and BRCA2 mutation carriers.

<sup>47</sup> 1-3 years of use.

<sup>48</sup> >3 years of use.

<sup>49</sup> 8 studies on BRCA1+ or BRCA2+ patients and breast cancer risk with CSC use.

## Breast Cancer References

- Adams-Campbell LL, Makambi KH, Frederick WA, Gaskins M, Dewitty RL, and McCaskill-Stevens W. Breast cancer risk assessments comparing Gail and CARE models in African-American women. *Breast J*. 15 Suppl 2009; 1:S72–75.
- Amadou A, Fabre A, Torres-Mejía G, Ortega-Olvera C, Angeles-Llerenas A, McKenzie F, Biessy C, Hainaut P, and Romieu I. Hormonal therapy and risk of breast cancer in Mexican women. *PLoS One* 2013; 8:e79695.
- Beaber EF, Buist DS, Barlow WE, Malone KE, Reed SD, and Li CI. Recent oral contraceptive use by formulation and breast cancer risk among women 20 to 49 years of age. *Cancer Res* 2014a; 74:4078–4089.
- Beaber EF, Malone KE, Tang MT, Barlow WE, Porter PL, Daling JR, and Li CI. Oral contraceptives and breast cancer risk overall and by molecular subtype among young women. *Cancer Epidemiol Biomarkers Prev* 2014b; 23:755–764.
- Beji NK and Reis N. Risk factors for breast cancer in Turkish women: a hospital-based case-control study. *Eur J Cancer Care (Engl)*. 2007; 16:178–184.
- Bethea TN, Rosenberg L, Hong CC, Troester MA, Lunetta KL, Bandera EV, Schedin P, Kolonel LN, Olshan AF, Ambrosone CB, and Palmer JR. A case-control analysis of oral contraceptive use and breast cancer subtypes in the African American Breast Cancer Epidemiology and Risk Consortium. *Breast Cancer Res* 2015; 17:22. Doi: 10.1186/s13058-015-0535-x.
- Brohet RM, Goldgar DE, Easton DF, Antoniou AC, Andrieu N, Chang-Claude J, Peock S, Eeles RA, Cook M, Chu C, Noguès C, Lasset C, Berthet P, Meijers-Heijboer H, Gerdes AM, Olsson H, Caldes T, van Leeuwen FE, and Rookus MA. Oral contraceptives and breast cancer risk in the international BRCA1/2 carrier cohort study: a report from EMBRACE, GENEPSO, GEO-HEBON, and the IBCCS Collaborating Group. *J Clin Oncol* 2007; 25:3831–3836.
- Delort L, Kwiatkowski F, Chalabi N, Satih S, Bignon YJ, and Bernard-Gallon DJ. Risk factors for early age at breast cancer onset—the “COSA program” population-based study. *Anticancer Res* 2007; 27:1087–1094.
- Dolle JM, Daling JR, White E, Brinton LA, Doody DR, Porter PL, and Malone KE. Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer Epidemiol Biomarkers Prev* 2009; 18:1157–1166.
- Figueiredo JC, Haile RW, Bernstein L, Malone KE, Largent J, Langholz B, Lynch CF, Bertelsen L, Capanu M, Concannon P, Borg A, Børresen-Dale AL, Diep A, Teraoka S, Torngren T, Xue S, and Bernstein JL. Oral contraceptives and postmenopausal hormones and risk of contralateral breast cancer among BRCA1 and BRCA2 mutation carriers and noncarriers: the WECARE Study. *Breast Cancer Res Treat* 2010; 120:175–183.
- Folger SG, Marchbanks PA, McDonald JA, Bernstein L, Ursin G, Berlin JA, Daling JR, Norman SA, Strom BL, Weiss LK, Simon MS, Burkman RT, Malone KE, and Spirtas R. Risk of breast cancer associated with short-term use of oral contraceptives. *Cancer Causes Control* 2007; 18:189–198.
- Friebel TM, Domchek SM, and Rebbeck TR. Modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: systematic review and meta-analysis. *J Natl Cancer Inst* 2014; 106(6):dju091. Doi: 10.1093/jnci/dju091.

Haile RW, Thomas DC, McGuire V, Felberg A, John EM, Milne RL, Hopper JL, Jenkins MA, Levine AJ, Daly MM, Buys SS, Senie RT, Andrulis IL, Knight JA, Godwin AK, Southey M, McCredie MR, Giles GG, Andrews L, Tucker K, Miron A, Apicella C, Tesoriero A, Bane A, Pike MC; kConFab Investigators; Ontario Cancer Genetics Network Investigators, and Whittemore AS. BRCA1 and BRCA2 mutation carriers, oral contraceptive use, and breast cancer before age 50. *Cancer Epidemiol Biomarkers Prev* 2006; 15:1863–1870.

Heikkinen S, Koskenvuo M, Malila N, Sarkeala T, Pukkala E, and Pitkäniemi J. Use of exogenous hormones and the risk of breast cancer: results from self-reported survey data with validity assessment. *Cancer Causes Control* 2016; 27:249–258.

Hunter DJ, Colditz GA, Hankinson SE, Malspeis S, Spiegelman D, Chen W, Stampfer MJ, and Willett WC. Oral contraceptive use and breast cancer: a prospective study of young women. *Cancer Epidemiol Biomarkers Prev* 2010; 19:2496–2502.

Ichida M, Kataoka A, Tsushima R, and Taguchi T. No increase in breast cancer risk in Japanese women taking oral contraceptives: a case-control study investigating reproductive, menstrual and familial risk factors for breast cancer. *Asian Pac J Cancer Prev* 2015; 16:3685–3690.

Jernström H, Loman N, Johannsson OT, Borg A, and Olsson H. Impact of teenage oral contraceptive use in a population-based series of early-onset breast cancer cases who have undergone BRCA mutation testing. *Eur J Cancer* 2005; 41:2312–2320.

Kahlenborn C, Modugno F, Potter DM, and Severs WB. Oral Contraceptive Use as a Risk Factor for Premenopausal Breast Cancer: A Meta-analysis. *Mayo Clin Proc* 2006; 81:1290–1302.

Kotsopoulos J, Lubinski J, Moller P, Lynch HT, Singer CF, Eng C, Neuhausen SL, Karlan B, Kim-Sing C, Huzarski T, Gronwald J, McCuaig J, Senter L, Tung N, Ghadirian P, Eisen A, Gilchrist D, Blum JL, Zakalik D, Pal T, Sun P, and Narod SA; Hereditary Breast Cancer Clinical Study Group. Timing of oral contraceptive use and the risk of breast cancer in BRCA1 mutation carriers. *Breast Cancer Res Treat* 2014; 143:579–586.

Lee E, Ma H, McKean-Cowdin R, Van Den Berg D, Bernstein L, Henderson BE, and Ursin G. Effect of reproductive factors and oral contraceptives on breast cancer risk in BRCA1/2 mutation carriers and noncarriers: results from a population-based study. *Cancer Epidemiol Biomarkers Prev* 2008; 17:3170–3178.

Li CI, Beaber EF, Tang MT, Porter PL, Daling JR, and Malone KE. Effect of depo-medroxyprogesterone acetate on breast cancer risk among women 20 to 44 years of age. *Cancer Res* 2012; 72:2028–2035.

Lumachi F, Frigo AC, Basso U, Tombolan V, and Ermani M. Estrogen therapy and risk of breast cancer in postmenopausal women: a case-control study and results of a multivariate analysis. *Menopause* 2010; 17:524–528.

Lund E, Bakken K, Dumeaux V, Andersen V, and Kumle M. Hormone replacement therapy and breast cancer in former users of oral contraceptives--The Norwegian Women and Cancer study. *Int J Cancer* 2007; 121:645–648.

Ma H, Bernstein L, Ross RK, and Ursin G. Hormone-related risk factors for breast cancer in women under age 50 years by estrogen and progesterone receptor status: results from a case-control and a case-case comparison. *Breast Cancer Res* 2006; 8:R39.

Ma H, Wang Y, Sullivan-Halley J, Weiss L, Marchbanks PA, Spirtas R, Ursin G, Burkman RT, Simon MS, Malone KE, Strom BL, McDonald JA, Press MF, and Bernstein L. Use of four biomarkers to evaluate the risk of breast cancer subtypes in the women's contraceptive and reproductive experiences study. *Cancer Res* 2010; 70:575–587.

Mehrgou A and Akouchekian M. The importance of BRCA1 and BRCA2 genes mutations in breast cancer development. *Med J Islam Repub Iran* 2016; 30:369.

Milne RL, Knight JA, John EM, Dite GS, Balbuena R, Ziogas A, Andrulis IL, West DW, Li FP, Southey MC, Giles GG, McCredie MR, Hopper JL, and Whittemore AS. Oral contraceptive use and risk of early-onset breast cancer in carriers and noncarriers of BRCA1 and BRCA2 mutations. *Cancer Epidemiol Biomarkers Prev* 2005; 14:350–356.

Moorman PG, Havrilesky LJ, Gierisch JM, Coeytaux RR, Lowery WJ, Peragallo Urrutia R, Dinan M, McBroom AJ, Hasselblad V, Sanders GD, and Myers ER. Oral contraceptives and risk of ovarian cancer and breast cancer among high-risk women: a systematic review and meta-analysis. *J Clin Oncol* 2013;31(33): 4188–4198. Doi: 10.1200/JCO.2013.48.9021.

Mørch LS, Skovlund CW, Hannaford PC, Iversen L, Fielding S, and Lidegaard Ø. Contemporary hormonal contraception and the risk of breast cancer. *N Engl J Med* 2017; 377:2228–2239.

Ozmen V, Ozcinar B, Karanlik H, Cabioglu N, Tukenmez M, Disci R, Ozmen T, Igci A, Muslumanoglu M, Kecer M, and Soran A. Breast cancer risk factors in Turkish women—a University Hospital based nested case control study. *World J Surg Oncol* 2009; 7:37.

Parkin DM. Cancers attributable to exposure to hormones in the UK in 2010. *British Journal of Cancer* 2011; 105: S42–S48.

Phipps AI, Chlebowski RT, Prentice R, McTiernan A, Wactawski-Wende J, Kuller LH, Adams-Campbell LL, Lane D, Stefanick ML, Vitolins M, Kabat GC, Rohan TE, and Li CI. Reproductive history and oral contraceptive use in relation to risk of triple-negative breast cancer. *J Natl Cancer Inst* 2011; 103:470–477.

Poosari A, Promthet S, Kamsa-ard S, Suwanrungruang K, Longkul J, and Wiangnon S. Hormonal contraceptive use and breast cancer in Thai women. *J Epidemiol* 2014; 24:216–220.

Rosenberg L, Zhang Y, Coogan PF, Strom BL, and Palmer JR. A case-control study of oral contraceptive use and incident breast cancer. *Am J Epidemiol* 2008; 169:473–479.

Rosenberg L, Boggs DA, Wise LA, Adams-Campbell LL, and Palmer JR. Oral contraceptive use and estrogen/progesterone receptor-negative breast cancer among African American women. *Cancer Epidemiol Biomarkers Prev* 2010; 19:2073–2079.

Samson ME, Adams SA, Mulatya CM, Zhang J, Bennett CL, Hebert J, and Steck SE. Types of oral contraceptives and breast cancer survival among women enrolled in Medicaid: A competing-risk model. *Maturitas* 2017; 95:42–49.

Silvera SA, Miller AB, and Rohan TE. Oral contraceptive use and risk of breast cancer among women with a family history of breast cancer: a prospective cohort study. *Cancer Causes Control*. 2005; 16:1059–1063.

Soini T, Hurskainen R, Grénman S, Mäenpää J, Paavonen J, and Pukkala E. Cancer risk in women using the levonorgestrel-releasing intrauterine system in Finland. *Obstet Gynecol* 2014; 124 (2 Pt 1): 292–299.

Sweeney C, Giuliano AR, Baumgartner KB, Byers T, Herrick JS, Edwards SL, and Slattery ML. Oral, injected and implanted contraceptives and breast cancer risk among U.S. Hispanic and non-Hispanic white women. *Int J Cancer* 2007; 121:2517–2523.

Thorbjarnardottir T, Olafsdottir EJ, Valdimarsdottir UA, Olafsson O, and Tryggvadottir L. Oral contraceptives, hormone replacement therapy and breast cancer risk: a cohort study of 16 928 women 48 years and older. *Acta Oncol* 2014; 53:752–758.

Trivers KF, Gammon MD, Abrahamson PE, Lund MJ, Flagg EW, Moorman PG, Kaufman JS, Cai J, Porter PL, Brinton LA, Eley JW, and Coates RJ. Oral contraceptives and survival in breast cancer patients aged 20 to 54 years. *Cancer Epidemiol Biomarkers Prev* 2007; 16:1822–1827.

Veisy A, Lotfinejad S, Salehi K, and Zhian F. Risk of breast cancer in relation to reproductive factors in North-West of Iran, 2013-2014. *Asian Pac J Cancer Prev* 2015; 16:451–455.

Veneroso C, Siegel R, and Levine PH. Early age at first childbirth associated with advanced tumor grade in breast cancer. *Cancer Detect Prev* 2008; 32:215–223.

Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting observational studies. *PLoS Medicine* 2007; 4:1623–27 (e296). Doi: 10.1371/journal.pmed. 0040296.

Williams WV, Carlson K, Mitchell LA, and Raviele K. Association of Combined Estrogen-Progestogen and Progestogen Only Contraceptives with the Development of Cancer. *The Linacre Quarterly* 2018; 85(4): 412–452.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives and Combined Estrogen–Progestogen Menopausal Therapy. 2007; <http://monographs.iarc.fr/ENG/Monographs/vol91/mono91.pdf>.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives. *IARC Monographs* 2012; 100A:283–318; <http://monographs.iarc.fr/ENG/Monographs/vol100A/mono100A-19.pdf>.

Yager JD and Davidson NE. Estrogen Carcinogenesis in Breast Cancer. *N Engl J Med* 2006; 354:270–282.

Zhu H, Lei X, Feng J, and Wang Y. Oral contraceptive use and risk of breast cancer: a meta-analysis of prospective cohort studies. *Eur J Contracept Reprod Health Care* 2012; 17:402–414.

## Cervical Cancer

According to the SEER statistics<sup>1</sup>, it is estimated that there are 257,524 women in the US with cervical cancer. There will be about 13,000 new cases of cervical cancer in 2018, with about 4,000 deaths. The five-year survival for cervical cancer is 66%. The IARC evaluation of an increased risk of cervical cancer with COCs is also supported especially by a large, high-quality cohort study (Roura 2016, Table 6). The data for cervical cancer presented in Table 4 shows in particular a higher risk for invasive cervical cancer, and a higher risk with current use. All studies appear to agree that there is an increased risk of cervical cancer in users of COCs (OR apparently about 1.05 per year of use), and this risk increases with duration of use. Current use appears to confer a higher risk than past use, and the risk for invasive cancer shows the highest increase in risk (Roura 2016). A meta-analysis of case-control studies that focused on patients positive for human papilloma virus DNA (Moreno 2002) also showed an increased risk, especially with protracted (5+ years) of use of COCs. One case-control study (McFarlane-Anderson 2008) and one meta-analysis (International Collaboration 2007) also showed an increased risk with progestogen-only contraceptives. Thus, there does appear to be an increased risk of cervical cancer in users of COCs or POCs, and the risk appears to increase with duration of use.

---

<sup>1</sup> <https://seer.cancer.gov/statfacts/html/corp.html>

Table 6 – Cervical Cancer

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Roura et al. 2016	Cohort Study		1.1 <sup>1</sup> (0.9–1.3)		1.8 <sup>10</sup> (1.4–2.4)		1 <sup>10</sup> (0.9–1.3)	1,065	306,971	94%
			1.6 <sup>2</sup> (1.1–2.3)		2.2 <sup>8</sup> (1.3–4.0)		1.6 <sup>8</sup> (1.1–2.2)	261	306,971	
Leslie et al. 2014	Case Control Study	1.35 <sup>3</sup> (0.99-1.85)						219	2,300	87%
McFarlane-Anderson et al. 2008	Case Control Study	1.59 <sup>4</sup> (0.87-2.82)						240	102	83%
		2.48 <sup>5</sup> (1.30-4.74)								
Vanakankovit et al. 2008	Case Control Study	1.49 (0.79-2.64)						60	180	76%
Wilson et al. 2013	Case Control Study	1.22 (0.96–1.56)						724	3,479	76%
Matos et al. 2005	Case Control Study	1.3 (0.8–3.1)						140	157	47%
International Collaboration 2007 <sup>6</sup>	Meta-analysis	1.057 (1.04–1.07)						16,573	35,509	97%
	<5 years of use	0.96 (0.04)8								
	5-9 years of use	1.2 (0.05)5								
	10+ years of use	1.56 (0.08)5								
	<5 years of use	1.07 (0.08)9						7,227	19,335	
	5+ years of use	1.22 (0.11)6								
Moreno 2002 <sup>10</sup>	Meta-analysis							1676	255	95%
	Invasive cervical cancer (ICC)	1.29 (0.88-1.91)								
	ICC 5+ years of use	4.01 (2.01-8.02)								

<sup>1</sup> Includes Cervical Intraepithelial Neoplasia Grade 3, carcinoma in situ and invasive cervical cancer.

<sup>2</sup> Analysis limited to invasive cervical cancer.

<sup>3</sup> Study limited to HIV+ women.

<sup>4</sup> Combined hormonal contraceptives.

<sup>5</sup> Progesterone only contraceptives.

<sup>6</sup> Meta-analysis of 24 studies (15 cohort and 9 case-control studies).

<sup>7</sup> Relative risk per year of use for current users of combined hormonal contraceptives.

<sup>8</sup> Floating standard error shown for users of combined hormonal contraceptives.

<sup>9</sup> Progestin only contraceptives. Floating standard error shown. The 95% CI for 5+ years of use is 1.01-1.46.

<sup>10</sup> Pooled data from 8 case-control studies of invasive cervical cancer and 2 of carcinoma in situ, analyzing only the subset positive for Human Papilloma Virus DNA in cervical cells.



Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
	In situ carcinoma (ISC)	1.42 (0.99-2.04)								
	ISC 5+ years of use	3.42 (2.13-5.48)								

## Cervical Cancer References

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives and Combined Estrogen–Progestogen Menopausal Therapy. 2007; <http://monographs.iarc.fr/ENG/Monographs/vol91/mono91.pdf>.

World Health Organization International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans. Combined Estrogen–Progestogen Contraceptives. *IARC Monographs* 2012; 100A:283–318; <http://monographs.iarc.fr/ENG/Monographs/vol100A/mono100A-19.pdf>.

International Collaboration of Epidemiological Studies of Cervical Cancer, Appleby P, Beral V, Berrington de González A, Colin D, Franceschi S, Goodhill A, Green J, Peto J, Plummer M, and Sweetland S. Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *Lancet* 2007; 370:1609–1621.

Leslie HH, Karasek DA, Harris LF, Chang E, Abdulrahim N, Maloba M, and Huchko MJ. Cervical cancer precursors and hormonal contraceptive use in HIV-positive women: application of a causal model and semi-parametric estimation methods. *PLoS One* 2014; 9:e101090.

Matos A, Moutinho J, Pinto D, and Medeiros R. The influence of smoking and other cofactors on the time to onset to cervical cancer in a southern European population. *Eur J Cancer Prev* 2005; 14:485–491.

McFarlane-Anderson N, Bazuaye PE, Jackson MD, Smikle M, and Fletcher HM. Cervical dysplasia and cancer and the use of hormonal contraceptives in Jamaican women. *BMC Womens Health* 2008; 8:9.

Moreno V, Bosch FX, Muñoz N, Meijer CJ, Shah KV, Walboomers JM, Herrero R, Franceschi S; International Agency for Research on Cancer. Multicentric Cervical Cancer Study Group. Effect of oral contraceptives on risk of cervical cancer in women with human papillomavirus infection: the IARC multicentric case-control study. *Lancet* 2002; 359(9312):1085–1092.

Roura E, Travier N, Waterboer T, de Sanjosé S, Bosch FX, Pawlita M, Pala V, Weiderpass E, Margall N, Dillner J, Gram IT, Tjønneland A, Munk C, Palli D, Khaw KT, Overvad K, Clavel-Chapelon F, Mesrine S, Fournier A, Fortner RT, Ose J, Steffen A, Trichopoulou A, Lagiou P, Orfanos P, Masala G, Tumino R, Sacerdote C, Polidoro S, Mattiello A, Lund E, Peeters PH, Bueno-de-Mesquita HB, Quirós JR, Sánchez MJ, Navarro C, Barricarte A, Larrañaga N, Ekström J, Lindquist D, Idahl A, Travis RC, Merritt MA, Gunter MJ, Rinaldi S, Tommasino M, Franceschi S, Riboli E, and Castellsagué X. The Influence of Hormonal Factors on the Risk of Developing Cervical Cancer and Pre-Cancer: Results from the EPIC Cohort. *PLoS One* 2016; 11:e0147029.

Vanakankovit N and Taneepanichskul S. Effect of oral contraceptives on risk of cervical cancer. *J Med Assoc Thai* 2008; 91:7–12.

Wilson JC, O'Rorke MA, Cooper JA, Murray LJ, Hughes CM, Gormley GJ, and Anderson LA. Non-steroidal anti-inflammatory drug use and cervical cancer risk: a case-control study using the Clinical Practice Research Datalink. *Cancer Epidemiol* 2013; 37:897–904.

## Crohn's Disease

Papers were accessed from a PubMed literature review as noted (Williams 2017). Each paper was rated based on the parameters noted in the STROBE statement (von Elm 2007).

Overall, 17 primary studies and two meta-analyses were identified which evaluated the effect of COCs on the later development of Crohn's disease (Table 7). Of the 17 primary studies, 4 showed a significantly increased risk for either ever use (Ng 2012, Sicilia 2001, Katschinski 1993) or current use (Katschinski 1993, Khalili 2013) or past use (Khalili 2013). None of the primary studies showed a significantly decreased risk. One meta-analysis (Godet 1995) gave a significantly increased RR of 1.44 (95% CI 1.12–1.86) for ever use of COCs. A meta-analysis published in 2008 showed a significantly increased risk for current use (RR of 1.46 [1.26–1.70]) compared with 1.04 (0.816–1.340) for past use. Recent studies have produced similar findings as older studies, with the highest OR published in 2012 (9.04 [1.11–73.6]). Overall these studies indicate that use of COCs conveys an increased risk of Crohn's disease, especially current use.

Table 7 – Individual Studies of the Effects of COCs on the Development of Crohn’s Disease

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Khalili et al. 2013 <sup>1</sup>	Cohort		1.43		2.82 (1.65–4.82)		1.39 (1.05–1.85)	315	117,060	93%
García Rodríguez et al. 2005 <sup>2</sup>	Cohort				1.94 (0.85–4.45)		1.04 (0.50–2.17)	171	10,000	88%
Logan and Kay 1989	Cohort		1.7 (0.88–3.2)					42	45,958	54%
Vessey et al. 1986 <sup>3</sup>	Cohort				1.33			18	17,014	46%
Boyko et al. 1994	Case-control		2 (1.0–3.7)					91	169	94%
Katschinski 1993 <sup>4</sup>	Case-control				2.5 (0.75–4.6)					93%
Katschinski 1993 <sup>5</sup>	Case-control				3.1 (1.1–6.7)					93%
Lashner et al. 1989	Case-control	1 (0.46–2.16)		0.73 (0.34–1.59)		1.8 (0.61–5.29)		51	51	88%
Lesko et al. 1985 <sup>6</sup>	Case-control		1.7 (1.0–3.2)					57	2189	83%
Sandler et al. 1992	Case-control		1.49 (0.99–2.26)					184	217	81%
Persson et al. 1993	Case-control		1.7 (0.9–3.2)					152	305	81%
Halfvarson et al. 2006 <sup>7</sup>	Case-control				1.5 (0.4–5.3)			102	102	75%
Lowe et al. 2009 <sup>8</sup>	Case-control		1.05					21,172	754,6131	74%
Ng et al. 2012 <sup>9</sup>	Case-control	4 (1.1–14.2)						125	125	74%

<sup>1</sup> Hazard ratios (RR adjusted for time).

<sup>2</sup> OR increased with duration of use.

<sup>3</sup> Authors’ calculation adjusted for smoking.

<sup>4</sup> Adjusted RR for 1-3 years prior to disease onset.

<sup>5</sup> Adjusted RR for >3 years prior to disease onset.

<sup>6</sup> RR is from multiple logistic regression analysis.

<sup>7</sup> Monozygotic and dizygotic twins.

<sup>8</sup> Adjusted incidence rate ratio.

<sup>9</sup> Twins study.

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Ng et al. 2012 <sup>10</sup>	Case-control	9.04 (1.11–73.6)								74%
Sicilia et al. 2001	Case-control	2.8 (1.01–7.77)						103	103	71%
Corrao et al. 1998	Case-control ever use			3.4 (1.0–11.9)		1.8 (0.4–7.3)		225	225	67%
Katschinski 1993 <sup>11</sup>	Case-control		4.3 (1.3–14.4)					83	83	57%
Han et al. 2010	Case-control		0.66 (0.38–1.15)					315	536	52%
Calkins et al. 1986 <sup>12</sup>	Case-control	1.14 (0.44–2.96)						66	67	42%
Calkins et al. 1986 <sup>13</sup>	Case-control	1.6 (0.59–4.37)						66	71	42%
Vcev et al. 2015	Case-control	0.28 (0.03–2.46)						11	42	31%
Cornish et al. 2008	Meta-analysis				1.46 (1.26–1.70)		1.04 (0.816–.340)	1251	74,564	91%
Cornish et al. 2008 <sup>14</sup>	Meta-analysis				1.58 (1.07–2.40)					91%
Godet et al. 1995 <sup>15</sup>	Meta-analysis		1.44 (1.12–1.86)					531	49,156	82%

<sup>10</sup> Multivariate analysis.

<sup>11</sup> RR for use >3 years.

<sup>12</sup> Hospital controls.

<sup>13</sup> Neighborhood controls.

<sup>14</sup> High quality studies.

<sup>15</sup> Adjusted for smoking.

## Crohn's Disease References

Boyko EJ, Theis MK, Vaughan TL, and Nicol-Blades B. Increased risk of inflammatory bowel disease associated with oral contraceptive use. *American Journal of Epidemiology* 1994; 140:268–278.

Calkins BM, Mendeloff AI, and Garland C. Inflammatory bowel disease in oral contraceptive users. *Gastroenterology* 1986; 91: 523–524.

Cornish JA, Tan E, Simillis C, Clark SK, Teare J, and Tekkis PP. The risk of oral contraceptives in the etiology of inflammatory bowel disease: a meta-analysis. *American Journal of Gastroenterology* 2008; 103:2394–2400.

Corrao G, Tragnone A, Caprilli R, Trallori G, Papi C, Andreoli A, Di Paolo M, Riegler G, Rigo GP, Ferraù O, Mansi C, Ingrosso M, and Valpiani D. Risk of inflammatory bowel disease attributable to smoking, oral contraception and breastfeeding in Italy: a nationwide case-control study. Cooperative Investigators of the Italian Group for the Study of the Colon and the Rectum (GISC). *International Journal of Epidemiology* 1998; 27:397–404.

García Rodríguez LA, González-Pérez A, Johansson S, and Wallander MA. Risk factors for inflammatory bowel disease in the general population. *Alimentary Pharmacology & Therapeutics* 2005; 22:309–315.

Godet PG, May GR, and Sutherland LR. Meta-analysis of the role of oral contraceptive agents in inflammatory bowel disease. *Gut* 1995; 37:668–673.

Halfvarson J, Jess T, Magnuson A, Montgomery SM, Orholm M, Tysk C, Binder V, and Järnerot G. Environmental factors in inflammatory bowel disease: a co-twin control study of a Swedish-Danish twin population. *Inflammatory Bowel Disease* 2006; 12:925–933.

Han DY, Fraser AG, Dryland P, and Ferguson LR. Environmental factors in the development of chronic inflammation: a case-control study on risk factors for Crohn's disease within New Zealand. *Mutation Research* 2010; 690:116–122.

Katschinski B. [Smoking and ovulation inhibitor in inflammatory bowel diseases]. *Medizinische Klinik (Munich, Germany)* 88 Suppl 1993; 1:5–8.

Khalili H, Higuchi LM, Ananthakrishnan AN, Richter JM, Feskanich D, Fuchs CS, and Chan AT. Oral contraceptives, reproductive factors and risk of inflammatory bowel disease. *Gut* 2013; 62:1153–1159.

Lashner BA, Kane SV, and Hanauer SB. Lack of association between oral contraceptive use and Crohn's disease: a community-based matched case-control study. *Gastroenterology* 1989; 97:1442–1447.

Lesko SM, Kaufman DW, Rosenberg L, Helmrich SP, Miller DR, Stolley PD, and Shapiro S. Evidence for an increased risk of Crohn's disease in oral contraceptive users. *Gastroenterology* 1985; 89:1046–1049.

Logan RF and Kay CR. Oral contraception, smoking and inflammatory bowel disease—findings in the Royal College of General Practitioners Oral Contraception Study. *International Journal of Epidemiology* 1989; 18: 105–107.

Lowe AM, Roy PO, Poulin M, Michel P, Bitton A, St-Onge L, and Brassard P. Epidemiology of Crohn's disease in Quebec, Canada. *Inflammatory Bowel Disease* 2009; 15:429–435.

- Ng SC, Woodrow S, Patel N, Subhani J, and Harbord M. Role of genetic and environmental factors in British twins with inflammatory bowel disease. *Inflammatory Bowel Disease* 2012; 18:725–736.
- Persson PG, Leijonmarck CE, Bernell O, Hellers G, and Ahlbom A. Risk indicators for inflammatory bowel disease. *International Journal of Epidemiology* 1993; 2(2):268–272.
- Sandler RS, Wurzelmann JI, and Lyles CM. Oral contraceptive use and the risk of inflammatory bowel disease. *Epidemiology* 1992; 3:374–378.
- Vcev A, Pezerovic D, Jovanovic Z, Nakic D, Vcev I, and Majnarić L. A retrospective, case-control study on traditional environmental risk factors in inflammatory bowel disease in Vukovar-Srijem County, north-eastern Croatia 2010. *Wiener Klinische Wochenschrift* 2015; 127:345–354.
- Vessey M, Jewell D, Smith A, Yeates D, and McPherson K. Chronic inflammatory bowel disease, cigarette smoking, and use of oral contraceptives: findings in a large cohort study of women of childbearing age. *British Medical Journal (Clinical Research Edition)* 1986; 292:1101–1113.

## Ulcerative Colitis

Papers were accessed from a PubMed literature review as noted (Williams 2017). Each paper was rated based on the parameters noted in the STROBE statement (von Elm 2007).

Overall 14 primary studies and one meta-analysis were identified which evaluated the effect of COCs on the later development of ulcerative colitis (Table 8). None of the primary studies has shown a statistically significant decrease in risk, while two showed a significant increase in risk for the development of ulcerative colitis with ever use of COCs (Boyko 1994, Parrello 1997). One meta-analysis examined ever use and failed to show a significant difference (Godet et al. 1995), while another meta-analysis examined current use and found a significantly increased relative risk of 1.28 (1.06–1.54). Overall these studies suggest that use of COCs conveys an increased risk of ulcerative colitis, especially current use.



Table 8 – Individual Studies of the Effects of COCs on the Development of Ulcerative Colitis

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Khalili et al. 2013 <sup>1</sup>	Cohort		1.18 (0.92–1.52)		1.22 (0.74–2.07)		1.18 (0.91–1.52)	392	116,983	93%
García Rodríguez et al. 2005	Cohort				1.58 (0.71–3.52)		0.67 (0.32–1.39)	222	10,000	88%
Logan and Kay 1989	Cohort		1.3 (0.82–2.0)					78	45,922	54%
Vessey et al. 1986 <sup>2</sup>	Cohort				2.1			31	17,001	46%
Boyko et al 1994	Case-control		1.7 (1.1–2.7)					211	341	94%
Lashner et al. 1990	Case-control	0.86 (0.40–1.85)		0.7 (0.27–1.83)		1.14 (0.41–1.15)		46	46	81%
Sandler et al. 1992 <sup>3</sup>	Case-control		1.1 (0.65–1.85)					89	217	81%
Persson et al. 1993	Case-control		1.7 (0.8–3.3)					145	305	81%
Halfvarson et al. 2006 <sup>4</sup>	Case-control				0.6 (0.1–2.5)			125	125	75%
Ng et al. 2012 <sup>5</sup>	Case-control	0.43 (0.11–1.66)						125	125	74%
Parrello et al. 1997 <sup>6</sup>	Case-control	3.11 (1.54–6.3)						536	755	67%
Corrao et al. 1998	Case-control			1.6 (0.9–3.0)		1.3 (0.6–2.8)		594	594	67%
Calkins et al. 1986 <sup>7</sup>	Case-control	0.62 (0.11–3.42)						35	32	42%
Calkins et al. 1986 <sup>8</sup>	Case-control	0.57 (0.11–2.88)						35	38	42%

<sup>1</sup> Hazard ratios (RR adjusted for time).

<sup>2</sup> Authors' calculation, adjusted for smoking.

<sup>3</sup> Interaction with smoking notes, higher RR in smokers (2.49).

<sup>4</sup> Monozygotic and dizygotic twins.

<sup>5</sup> Twins studies.

<sup>6</sup> Unclear how the calculation was done.

<sup>7</sup> Hospital controls.

<sup>8</sup> Neighborhood controls.

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Vcev et al. 2015	Case-control	0.75 (0.30–1.88)						62	42	31%
Cornish et al. 2008	Meta-analysis				1.28 (1.06–1.54)		1.07 (0.702–1.640)	883	74,932	91%
Cornish et al. 2008 <sup>9</sup>	Meta-analysis				1.24 (0.999–1.54)					91%
Godet et al. 1995 <sup>10</sup>	Meta-analysis		1.29 (0.94–1.77)					851	49,875	82%

---

<sup>9</sup> High quality studies.

<sup>10</sup> Adjusted for smoking.

## Ulcerative Colitis References

Boyko EJ, Theis MK, Vaughan TL, and Nicol-Blades B. Increased risk of inflammatory bowel disease associated with oral contraceptive use. *American Journal of Epidemiology* 1994; 140:268–278.

Calkins BM, Mendeloff AI, and Garland C. Inflammatory bowel disease in oral contraceptive users. *Gastroenterology* 1986; 91:523–524.

Cornish JA, Tan E, Simillis C, Clark SK, Teare J, and Tekkis PP. The risk of oral contraceptives in the etiology of inflammatory bowel disease: a meta-analysis. *American Journal of Gastroenterology* 2008; 103:2394–2400.

Corrao G, Tragnone A, Caprilli R, Trallori G, Papi C, Andreoli A, Di Paolo M, Riegler G, Rigo GP, Ferraù O, Mansi C, Ingrosso M, and Valpiani D. Risk of inflammatory bowel disease attributable to smoking, oral contraception and breastfeeding in Italy: a nationwide case-control study. Cooperative Investigators of the Italian Group for the Study of the Colon and the Rectum (GISC). *International Journal of Epidemiology* 1998; 27:397–404.

García Rodríguez LA, González-Pérez A, Johansson S, and Wallander MA. Risk factors for inflammatory bowel disease in the general population. *Alimentary Pharmacology & Therapeutics* 2005; 22:309–315.

Godet PG, May GR, and Sutherland LR. Meta-analysis of the role of oral contraceptive agents in inflammatory bowel disease. *Gut* 1995; 37:668–673.

Halfvarson J, Jess T, Magnuson A, Montgomery SM, Orholm M, Tysk C, Binder V, and Järnerot G. Environmental factors in inflammatory bowel disease: a co-twin control study of a Swedish-Danish twin population. *Inflammatory Bowel Disease* 2006; 12:925–933.

Khalili H, Higuchi LM, Ananthakrishnan AN, Richter JM, Feskanich D, Fuchs CS, and Chan AT. Oral contraceptives, reproductive factors and risk of inflammatory bowel disease. *Gut* 2013; 62:1153–1159.

Lashner BA, Kane SV, and Hanauer SB. Lack of association between oral contraceptive use and ulcerative colitis. *Gastroenterology* 1990; 99:1032–36.

Logan RF and Kay CR. Oral contraception, smoking and inflammatory bowel disease—findings in the Royal College of General Practitioners Oral Contraception Study. *International Journal of Epidemiology* 1989;18:105–107.

Ng SC, Woodrow S, Patel N, Subhani J, and Harbord M. Role of genetic and environmental factors in British twins with inflammatory bowel disease. *Inflammatory Bowel Disease* 2012; 18:725–736.

Parrello T, Pavia M, Angelillo IF, Monteleone G, Riegler G, Papi G, D'Inca R, Annese V, Tonelli F, Caprilli R, and Pallone F. Appendectomy is an independent protective factor for ulcerative colitis: results of a multicentre case control study. The Italian Group for the Study of the Colon and Rectum (GISC). *Italian Journal of Gastroenterology and Hepatology* 1997; 29:208–211.

Persson PG, Leijonmarck CE, Bernell O, Hellers G, and Ahlbom A. Risk indicators for inflammatory bowel disease. *International Journal of Epidemiology* 1993; 22:268–272.

Sandler RS, Wurzelmann JI, and Lyles CM. Oral contraceptive use and the risk of inflammatory bowel disease. *Epidemiology* 1992; 3:374–378.

Vcev A, Pezerovic D, Jovanovic Z, Nakic D, Vcev I, and Majnarić L. A retrospective, case-control study on traditional environmental risk factors in inflammatory bowel disease in Vukovar-Srijem County, north-eastern Croatia, 2010. *Wiener Klinische Wochenschrift* 2015; 127:345–354.

Vessey M, Jewell D, Smith A, Yeates D, and McPherson K. Chronic inflammatory bowel disease, cigarette smoking, and use of oral contraceptives: findings in a large cohort study of women of childbearing age. *British Medical Journal (Clinical Research Edition)* 1986; 292:1101–1113.

## Systemic Lupus Erythematosus

Papers were accessed from a PubMed literature review as noted (Williams 2017). Each paper was rated based on the parameters noted in the STROBE statement (von Elm 2007).

There have been seven studies published evaluating the effect of hormonal contraceptives on susceptibility to systemic lupus erythematosus (Table 9). A significantly increased risk for development of systemic lupus erythematosus with use of COCs was shown for ever use in two studies (Costenbader 2007, Sanchez-Guerrero 1997), for current use in one study (Bernier 2009) and for past use in one study (Costenbader 2007). None of the studies showed a decreased risk. While no meta-analyses of these studies have been performed, the uniformity of the results implicate COCs as an important risk factor for the subsequent development of systemic lupus erythematosus.

Table 9 – Individual Studies of the Effects of COCs on the Development of Systemic Lupus Erythematosus

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Costenbader et al. 2007 <sup>1</sup>	Cohort		1.5 (1.1–2.1)				1.7 (1.2-2.3)	262	238,046	96%
Costenbader et al. 2007 <sup>2</sup>	Cohort		1.6 (1.1-2.2)				1.6 (1.1-2.2)	164	102,882	96%
Costenbader et al. 2007 <sup>3</sup>	Cohort		2.3 (1.0-5.0)				2.3 (1.1-5.2)	98	107,854	96%
Bernier et al. 2009	Cohort		1.19 (0.98-1.45)		1.54 (1.15-2.07)		1.06 (0.85-1.33)	786	7817	96%
Bernier et al. 2009 <sup>4</sup>	Cohort				2.52 (1.14-5.57)			786	7817	96%
Bernier et al. 2009 <sup>5</sup>	Cohort				1.45 (1.06-1.99)			786	7817	96%
Sanchez-Guerrero et al. 1997	Cohort		1.4 (0.9-2.1)					99	121,546	88%
Sanchez-Guerrero et al. 1997 <sup>6</sup>	Cohort		1.9 (1.1-3.3)					58	121,587	88%
Cooper et al. 2002	Case-control			1.5 (0.8–2.7)		1.3 (0.8–2.0)		240	321	92%
Strom et al. 1994	Case-control	0.8 (0.5-1.4)						195	143	73%
Zonana-Nacach et al. 2002 <sup>7</sup>	Case-control	2.1 (1.18-3.6)						130	130	61%
Grimes et al. 1985	Case-control			0.5 (0.11-2.3)				109	109	58%

<sup>1</sup> Pooled RR from the Nurses' Health Study (NHS) and NHS II.

<sup>2</sup> RR from the NHS (data collection through 1976).

<sup>3</sup> RR from NHS II (data collection through 1989).

<sup>4</sup> RR for short term use (starting COCs within ≤3 months).

<sup>5</sup> RR for long term use (starting COCs over 3 months previously with current use ongoing).

<sup>6</sup> Using most stringent definition of systemic lupus erythematosus.

<sup>7</sup> Paper written in Spanish. OR is for use of oral contraceptives for more than one year.

## Systemic Lupus Erythematosus References

Bernier MO, Mikaeloff Y, Hudson M, and Suissa S. Combined oral contraceptive use and the risk of systemic lupus erythematosus. *Arthritis and Rheumatism* 2009; 61:476–481.

Cooper GS, Dooley MA, Treadwell EL, St Clair EW, and Gilkeson GS. Hormonal and reproductive risk factors for development of systemic lupus erythematosus: results of a population-based, case-control study. *Arthritis and Rheumatism* 2002; 46:1830–1839.

Costenbader KH, Feskanich D, Stampfer MJ, and Karlson EW. Reproductive and menopausal factors and risk of systemic lupus erythematosus in women. *Arthritis and Rheumatism* 2007; 56:1251–1262.

Grimes DA, LeBolt SA, Grimes KR, and Wingo PA. Systemic lupus erythematosus and reproductive function: A case control study. *American Journal of Obstetrics and Gynecology* 1985; 153:179–186.

Sanchez-Guerrero J, Karlson EW, Liang MH, Hunter DJ, Speizer FE, and Colditz GA. Past use of oral contraceptives and the risk of developing systemic lupus erythematosus. *Arthritis and Rheumatism* 1997; 40: 804–808.

Strom BL, Reidenberg MM, West S, Snyder ES, Freundlich B, and Stolley PD. Shingles, allergies, family medical history, oral contraceptives, and other potential risk factors for systemic lupus erythematosus. *American Journal of Epidemiology* 1994; 140:632–642.

Zonana-Nacach A, Rodríguez-Guzmán LM, Jiménez-Balderas FJ, Camargo-Coronel A, Escobedo-de la Peña J, and Fraga A. [Risk factors associated with systemic lupus erythematosus in a Mexican population]. *Salud Pública de México* 2002; 44:213–218.

## Risk of Depression, Mood Disorders, and Suicide

The effects of contraceptive steroid hormones on depression, mood disorders, and suicide have been investigated (Table 10). The largest study of incident depression and use of anti-depressant medication (Skovlund 2016) indicates significantly increased risks for both COCs and POCs for both outcomes. The same group studied for suicide attempts and suicides (Skovlund 2018). Elevated risks were seen, and this was the case for both COCs and POCs. The recent NCHA study (Gregory 2018) showed a similar trend. One study (Keyes 2013) showed a lower risk of depression, but was not measuring clinically diagnosed depression, but rather the presence of depressive symptoms within 7 days prior to the survey. They also found a lower rate of suicide attempts among COC users. Similar findings were seen in 2 studies that also used a questionnaire looking at current COC or POC use (Toffol 2011, Toffol 2012). An analysis of the development of mood disorders found a higher incidence with POCs but a lower incidence with COCs (Svendal 2012). A study of post-partum depression as a reported adverse drug reaction showed higher rates for levonogestrel, etonogestrel and sertraline & drospirenone (Horibe 2018). A study of post-partum DMPA versus copper IUD use showed significant increases in depression scores and major depressive episodes with DMPA (Singata-Madliki, 2016). A retrospective cohort study showed increased risk for antidepressant use in patients who used ethinyl estradiol/etonogestrel (ring), and decreased risk of depression diagnosis with norethindrone-only pills or the levonorgestrel intrauterine system. A small retrospective chart review of the effect of immediate post-partum DMPA did not show significant effects on post-partum depression (Tsai 2009). All the papers, which have broken out the age groups of users, show maximum increased risk for depression, suicide risk, and suicide within 3 months of beginning to use the drugs and tapering off after 6 months, partly due to attenuation of symptoms, partly due to discontinuation due to adverse effects. These risks need to be adequately conveyed in prescribing information and patient-related materials.

However, little attention has been paid to the effects of blocking the important actions of estradiol and progesterone with progestins during the time of active brain remodeling. Estradiol and progesterone in normal sequence are essential for brain remodeling from ages 15–19 years particularly for myelination, dendritic pruning and establishment of new synaptic connections (Del Rio 2018). Suppressing these with synthetic progestins can have far-reaching, untoward effects. See Griksiene below in Table 10 as well as Del Rio (Del Rio 2018).



Table 10 – Studies of Chemical Contraceptives and Depression, Mood Disorders and Suicides

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
Skovlund 2016 incl /Worley	Prospective Cohort Incident Depression – COCs		1.1 <sup>95</sup> (1.08-1.14)				1,061,997
	Incident Depression – POCs		1.2 <sup>96</sup> (1.04-1.31)				
	First use of Antidepressants – COCs		1.23 <sup>97</sup> (1.22-1.25)				
	First use of Antidepressants – POCs		1.3 <sup>98</sup> (1.27-1.40)				
Skovlund 2018 incl /Worley	Prospective Cohort						475,802
	Prospective Cohort Suicide attempts		1.97 <sup>99</sup> (1.85-2.10)				
	Suicides		3.08 <sup>100</sup> (1.34-7.08)				
Gregory 2018	NCHA survey					146,938	202,759
	Ever Diagnosed with Depression	1.558 (1.506-1.612)					
	Academic performance affected by depression	1.282 (1.245-1.321)					
Keyes 2013	COC reduced depression among women 25-34 years of age. <sup>101</sup> 4 waves of L-Hanes			-1.04 <sup>102</sup> (-1.73 - -0.35)		3224	1219
	Suicide attempts			0.38 (0.15-0.97)			
Toffol 2011	Population/choice			-0.988 <sup>104</sup> (-1.917 – -0.059)			2,310

<sup>95</sup> First diagnosis of depression for combined oral contraceptive users.

<sup>96</sup> First diagnosis of depression for all progestin-only method users.

<sup>97</sup> First use of an antidepressant for combined oral contraceptive users.

<sup>98</sup> First use of an antidepressant for all progestin-only method users.

<sup>99</sup> Hazard ratio for suicide attempts; all hormonal contraceptives.

<sup>100</sup> Hazard ratio for suicides; all hormonal contraceptives.

<sup>101</sup> “The presence of depressive symptoms during the past 7 days was assessed in all waves using the Center for Epidemiologic Studies Depression Scale (CES-D).”

<sup>102</sup>  $\beta$  statistic shown.

<sup>104</sup>  $\beta$  statistic shown for the Beck Depression Inventory (BDI). None of the other parameters assessed was statistically significant (including any psychiatric diagnosis, alcohol dependence, major depressive episode or disorder, dysthymic disorder, or anxiety disorder).

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
	Cross sectional 30-54 yrs. of age <sup>103</sup>						
Toffol 2012	Population-based cross-sectional study <sup>105</sup>			-0.42 (-1.79 - -0.04) <sup>106</sup>			8,586
Svendal 2012 <sup>107</sup>	Population-based cross-sectional study					40	458
	POC Use – mood disorder			3.0 (1.1-7.8)			
	COC Use – mood disorder			0.3 (0.1-0.9)			
Horibe 2018	Retrospective <sup>108</sup>					253	6,157,897
	Post-partum depression w/ levonorgestrel			12.5 (8.7-18)			
	Post-partum depression w/ etonogestrel			14.0 (8.5-22.8)			
	Post-partum depression w/ sertraline & drospirenone			5.4 (2.7-10.9)			
Singata-Madliki 2016	Single-blind randomized controlled trial of post-partum DMPA vs. copper IUD			109		111 <sup>110</sup>	117 <sup>111</sup>

<sup>103</sup> “The associations between the current use of COCs and the LNG-IUS, and their duration versus mood symptoms [Beck Depression Inventory (BDI)], psychological well-being [(General Health Questionnaire-12 (GHQ-12))] and recent psychiatric diagnoses [(Composite International Diagnostic Interview (CIDI))] were examined among women who participated in the Finnish-population-based Health 2000 study.” “Overall, hormonal contraception was well tolerated with few significant effects on psychological well-being.”

<sup>105</sup> Data were collected in the context of the National FINRISK Study Survey, a cross-sectional population-based health survey carried out in Finland every 5 years since 1972. For the purpose of this study, data collected in the years 1997, 2002 and 2007 were analyzed for ages 25–54. OC vs. LNG. inconsistent questions between surveys, BDI, recall bias, etc. “Presence of somatic and psychological symptoms was assessed by asking the participants how often (often, sometimes, not at all) in the previous month they had had one or more out of 13 symptoms.” Also administered the Beck Depression Inventory-13. “A negative association between the current use of COCs and Beck Depression Inventory-13 (BDI-13) score was found. Some other negative associations, all characterized by a small effect size, were detected between current use of COCs and the BDI items feelings of dissatisfaction, feelings of uselessness, irritability, lost interest in people and lost appetite.”

<sup>106</sup> Results for the BDI-13 shown. Other parameters (including BDI-21, low mood last year, anhedonia last year, recent diagnosis of depression and recent other psychiatric diagnosis) did not reach statistical significance.

<sup>107</sup> Women in Australia 20-50 years of age. Evaluated for the occurrence of mood disorders, including major depressive disorder (MDD), minor depression, bipolar disorder, dysthymia, mood disorder due to a general medical condition and substance induced mood disorder.

<sup>108</sup> Data is from the FDA Adverse Event Reporting System (FAERS) database. Reporting Odds Ratios (ROR) are shown.

<sup>109</sup> Beck Depression Inventory (BDI-II) and the Edinburgh Postnatal Depression Scale (EPDS) evaluated. The one-month EPDS depression scores were statistically significantly higher in the DMPA arm compared with the IUD arm ( $p=0.04$ ). Three-month BDI-II scores were significantly higher in the DMPA arm than in the IUD arm ( $p=0.002$ ) and, according to the BDI-II but not the EPDS, more women in the DMPA arm had major depression at this time-point (8 vs 2;  $p=0.05$ ).

<sup>110</sup> 111 randomized to DMPA.

<sup>111</sup> 117 randomized to IUDs.

	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	Cases	Controls/Cohort Size
Kulkarni 2005 <sup>112</sup>	Case-control pilot study COCs vs non-users			p=0.001 depression for all scales <sup>113</sup>		26	32
Roberts 2017	Retrospective cohort study <sup>114</sup>			With Dx of depression <sup>115</sup>	w/anti depressant use <sup>116</sup>	31,506 <sup>117</sup>	44,022 <sup>118</sup>
	Norethindrone-only pills			0.56 (0.49-0.64)	0.58 (0.52-0.64)		
	Levonorgestrel intrauterine system			0.65 (0.52-0.82)	1.01 (0.87-1.18)		
	Etonogestrel subdermal implant			1.01 (0.83-1.22)	1.22 (1.06-1.41)		
	Ethinyl estradiol/ norgestimate (pill)			0.89 (0.70-1.14)	1.02 (0.85-1.22)		
	Ethinyl estradiol/norethindrone (pill)			0.82 (0.59-1.12)	0.88 (0.69-1.13)		
	Ethinyl estradiol/etonogestrel (ring)			1.09 (0.80-1.50)	1.45 (1.16-1.80)		
Tsai 2009	Retrospective chart review <sup>119</sup>	DMPA	Controls			55	192
	Mean EPDS scores at 6 weeks postpartum	5.02	6.17				
Griksiene 2011	Case-control study <sup>120</sup>	<sup>121</sup>				23 <sup>122</sup>	20 <sup>123</sup>

<sup>112</sup> Assessment tools included three depression rating scales: Beck Depression Inventory (BDI), Hamilton Rating Scale for Depression (HAM-D) and Montgomery-Asberg Depression Rating Scale (MADRS); also used the Global Assessment of Functioning (GAF) Scale.

<sup>113</sup> ANOVA of GAF, BDI, HAM-D & MADRS scales all significantly different.

<sup>114</sup> Post-partum depression with hormonal contraception.

<sup>115</sup> Adjusted hazard ratios shown.

<sup>116</sup> Adjusted hazard ratios shown.

<sup>117</sup> Number on hormonal contraceptives.

<sup>118</sup> Number not on hormonal contraceptives.

<sup>119</sup> Depot medroxyprogesterone in the immediate post-partum period and depression. Evaluated the Edinburgh Postnatal Depression Scale (EPDS).

<sup>120</sup> Verbal fluency and mental rotation (spatial perception) are affected by progestins w/androgenic or antiandrogenic properties.

<sup>121</sup> Naturally cycling women performed better on verbal fluency task as compared to OC users. Subjects who used the third generation (androgenic) COCs generated significantly fewer words as compared to new generation (anti-androgenic) OC users and non-users. The third generation OC users demonstrated significantly longer RT in MRT task as compared to non-users. The MRT, verbal fluency and mood parameters did not depend on the phase of menstrual cycle.

<sup>122</sup> Women on hormonal contraception.

<sup>123</sup> Control women not on hormonal contraception.

## Depression, Mood Disorders, and Suicide References

Del Rio JP, Allende MI, Molina N, Serrano FG, Molina S, and Vigil P. Steroid Hormones and their Action in Women's Brains: The Importance of Hormonal Balance. *Frontiers in Public Health* 2018; May(6) art. 141:1–15.

Gregory Sean T, Hall K, Quast T, Gatto A, Bleck J, Storch EA, and DeBate R. Hormonal contraception, depression and Academic Performance among females attending college in the United States. *Psychiatry Research* 2018; 270:111–116.

Keyes Katherine T, Cheslack-Postava K, Westhoff C, Heim CM, Haloosim M, Walsh K, and Koenen K. Association of Hormonal Contraception Use with Reduced levels of Depressive Symptoms: A National Study of Sexually Active women in the United States. *Am J. Epidemiol* 2013; 178(9):1378–1388.

Skovlund CW, Mørch LS, Kessling LV, and Lidegaard O. Association of Hormonal Contraception with Depression. *JAMA Psychiatry* 2016; 73(11):1154–1162.

Skovlund CW, Mørch LS, Kessling LV, Lange T, and Lidegaard, O. Association of Hormonal Contraception with Suicide Attempts and Suicides. *Am. J Psychiatry* 2018; 175(4):336–342.

Svendal G, Berk M, Pasco JA, and Jacka FN. The use of hormonal contraceptive agents and mood disorders in women. *J Affective Disorders* 2012; 140:92–96.

Toffol E, Heiknheimo, Koponene P, Luoto R, and Partonen T. Hormonal contraception and mental health: results of a population based study. *Human Reproduction* 2011; 26(11):3085–3093.

Worly Brett L, Gur TL, and Schaffir J. The relationship between progestin hormonal contraception and depression: a systematic review. *Contraception* 2018; 97:478–489.

Young EA, Kornstein SG, Harvey AT, Wisniewski SR, Barkin J, Fava M, Trivedi MH, and Rush AJ. Influences of Hormone-Based Contraception in Depressive symptoms in Premenopausal Women with Major Depression. *Psychoneuroendocrinology* 2007; 32(7):843–853.

## Multiple Sclerosis

Papers were accessed from a PubMed literature review as noted (Williams 2017). Each paper was rated based on the parameters noted in the STROBE statement (von Elm 2007).

A total of 6 studies (3 cohort studies and 3 case-control studies) were identified which evaluated the impact of COCs on the subsequent development of multiple sclerosis (Table 11). Two studies showed a significantly increased risk for the development of multiple sclerosis with ever use of COCs (Hellwig 2016, Kotzamani 2012) with a similarly increased risk noted in one study for current use or past use (Hellwig 2016). Overall these studies suggest that use of COCs may convey an increased risk for the subsequent development of multiple sclerosis.

Table 11 – Individual Studies of the Effects of COCs on the Development of Multiple Sclerosis

Study	Study Design	OR Ever Use	RR Ever Use	OR Current Use	RR Current Use	OR Past Use	RR Past Use	Cases	Controls	Quality Score
Hernán et al. 2000 <sup>124</sup>	Cohort		1.1 (0.9-1.5)		1 (0.6-1.6)		1.2 (0.9-1.5)	313	237,318	90%
Thorogood et al. 1998 <sup>125</sup>	Cohort				1.2 (0.7-2.0)		1.3 (0.9-2.0)	114	46,000	75%
Villard-Mackintosh et al. 1993	Cohort		0.8 (0.5-1.4)					63	16,969	65%
Hellwig et al. 2016	Case-control	1.51 (1.12-2.03)		1.47 1.05-2.05		1.55 (1.20-2.00)		400	3804	92%
Kotzamani et al. 2012	Case-control	1.6 (1.1-2.4)						254	314	81%
Alonso et al. 2005 <sup>126</sup>	Case-control	0.6 (0.4-1.0)		0.5 (0.3-1.2)		0.6 (0.4-1.0)		106	1001	77%

<sup>124</sup> NHS I and II cohorts.

<sup>125</sup> Funded by drug companies that make HCs.

<sup>126</sup> OC use over the 3 years prior to the index date. Limited to women ≤50 years of age.

## Multiple Sclerosis References

Alonso A, Jick SS, Olek MJ, Ascherio A, Jick H, and Hernán MA. Recent use of oral contraceptives and the risk of multiple sclerosis. *Archives of Neurology* 2005; 62:1362–1365.

Hellwig K, Chen LH, Stanczyk FZ, and Langer-Gould AM. Oral Contraceptives and Multiple Sclerosis/Clinically Isolated Syndrome Susceptibility. *PLoS One* 2016; 11:e0149094. Doi:10.1371/journal.pone.0149094.

Hernán MA, Hohol MJ, Olek MJ, Spiegelman D, and Ascherio A. Oral contraceptives and the incidence of multiple sclerosis. *Neurology* 2000; 55:848–854.

Kotzamani D, Panou T, Mastorodemos V, Tzagournissakis M, Nikolakaki H, Spanaki C, and Plaitakis A. Rising incidence of multiple sclerosis in females associated with urbanization. *Neurology* 2012; 78:1728–1735.

Thorogood M, and Hannaford PC. The influence of oral contraceptives on the risk of multiple sclerosis. *British Journal of Obstetrics and Gynaecology* 1998; 105:1296–1299.

Villard-Mackintosh L, and Vessey MP. Oral contraceptives and reproductive factors in multiple sclerosis incidence. *Contraception* 1993; 47:161–168.

## Interstitial Cystitis

A case-control study (Konkle 2012) showed significantly higher use of birth control pills in cases versus controls: 88% versus 82%;  $P = 0.019$ . Another case-control study showed that use of COCs markedly increased the risk of the disease whether past (OR 4.6, 95% CI 1.74-12.1) or current use (OR 6.9, 95% CI 2.1–22.1). Interstitial cystitis was associated with vulvodynia and sexual dysfunction in a high number of cases (Gardella 2011). Another study showed that use of COCs in patients with interstitial cystitis was associated with a decrease in quality of life (El Khoudary 2009). One meta-analysis (Champaneria 2015) showed that ever use of COCs significantly increased the risk of interstitial cystitis (OR 2.31, 95% CI 1.03–5.16).

Overall, use of COCs appears to be associated with an increased risk for the development of interstitial cystitis.

## Interstitial Cystitis References

Champaneria R, D'Andrea RM, and Latthe PM. Hormonal contraception and pelvic floor dysfunction: a systematic review. *Int Urogynecol J* 2016; 27:709–722.

El Khoudary SR, Talbott EO, Bromberger JT, Chang CC, Songer TJ, and Davis EL. Severity of interstitial cystitis symptoms and quality of life in female patients. *Journal of Womens Health (Larchmont)* 2009; 18:1361–1368. Doi: 10.1089/jwh.2008.1270.

Gardella B, Porru D, Nappi RE, Daccò MD, Chiesa A, and Spinillo A. Interstitial cystitis is associated with vulvodynia and sexual dysfunction—a case-control study. *The Journal of Sexual Medicine* 2011; 8:1726–1734. Doi: 10.1111/j.1743-6109.2011.02251.x. Epub 2011 Apr 7.

Konkle K, Berry SH, Elliott MN, Hilton L, Suttorp MJ, Clauw DJ, and Clemens JQ. Comparison of an interstitial cystitis/bladder pain syndrome clinical cohort with symptomatic community women from the RAND Interstitial Cystitis Epidemiology study. *Journal of Urology* 2012; 187:508–512.



## Osteoporotic Bone Fractures

Prescribing information for POCs typically includes a warning regarding the development of osteoporosis. However, the more relevant outcome is fracture risk. Therefore, articles were sought that looked at the effect of COCs and POCs on fracture risk. Data were initially derived from a systematic review of the evidence from observational studies of hormonal contraceptive use for contraception and the risk of fracture in women by Lopez (Lopez 2015). They noted that in 2004, the US Food and Drug Administration added a warning to depot medroxyprogesterone acetate (DMPA) labeling about the potential loss of BMD (FDA 2004), which might limit long-term use. A systematic review of progestin-only methods found an association between DMPA use and loss of bone mineral density (Curtis 2006). Lopez identified 559 records, 524 of which did not meet their inclusion criteria. Thirty-five full-text reports remained, 11 of which were excluded. Of the remaining 24, 10 were secondary articles. That left 14 articles: the 14 studies examined oral contraceptives (N = 12), DMPA (N = 4) and the hormonal IUD (N = 1). Similar search terms to Lopez were used for papers published since 2015 and 2 additional papers were retrieved. The resulting studies are shown in Table 12.

COCs: Three early studies (Cooper 1993, Tuppurainen 1993, Vessey 1998) showed an increase risk of fracture with use of COCs. These studies predominately evaluated pre-menopausal fracture risk. Others that evaluated wrist fracture linked to falling had few cases but showed a trend to decreased risk (O'Neill 1996). One study that evaluated post-menopausal fracture risk based on prior oral contraceptive use (Barad 2005) also found an increased fracture risk. Another study looking at hip fracture risk in elderly women (Michaëlsson 1999) showed a decreased risk but is compromised in that "The exposure time for oral contraceptives may thus maximally have spanned 5 years..." Two studies by Vestergaard (Vestergaard 2006 and Vestergaard 2008) looked at any fracture with OC use and did not show a significant effect when multivariate analyses were performed. However, these studies only looked at use within the past 5 years and did not take into account remote use or cumulative lifetime use. A small cross-sectional study in southern Tasmania (Wei 2011) was stratified by duration of use and showed a reduction in vertebral deformities for 5-10 years of use, but no effect for shorter or longer duration of use and no effect on number of vertebral deformities. A large case-control study which evaluated incident fracture risk with varying numbers of COC prescriptions showed an increased risk for 10+ prescriptions with current use (Meier 2010). A similar study failed to confirm this for most prescription numbers (Kyvernitakis 2016) but this study had fewer subjects reducing its power. A case-control study (Memon 2011) nested in an earlier cohort study (Cooper 1993) failed to show an effect.

Overall the weight of evidence for use of COCs suggests an increased risk of bone fracture with protracted use. The study by Barad (2005) appears to have the largest number of subjects, was a cohort study, and was the only study that evaluated post-menopausal fracture risk with prior use of COCs.

In contrast, virtually all the studies evaluating POCs show an elevated risk (Lanza 2013, Vestergaard 2008b, Meier 2010, Kyvernitakis 2016). This risk appears to increase with duration of use.

Table 12 – Individual Studies of the Effects of Contraceptives on the Development of Osteoporotic Fractures

Study	Study Design	Intervention	OR	RR	Cases	Controls or Cohort Size	Outcome
Cooper 1993 <sup>127</sup>	Cohort	COCs		1.20 (1.08-1.34)	1365	46,000	All fractures
Vessey 1998 <sup>128</sup>	Cohort	COCs		1.5 (1.1-2.1)	1308	17,032	First fracture: radius or ulna
Vessey 1998 <sup>129</sup>	Cohort	COCs		1.2 (1.1-1.4)			First fracture: all sites
Vessey 1998 <sup>130</sup>	Cohort	COCs		2.5 (1.5-4.0)			First fracture: radius or ulna
Vessey 1998 <sup>131</sup>	Cohort	COCs		1.3 (1.1-1.5)			First fracture: all sites
Vessey 1998 <sup>132</sup>	Cohort	COCs		5.7 (p=0.017)			First fracture: radius or ulna
Vessey 1998 <sup>133</sup>	Cohort	COCs		11.2 (p<0.001)			First fracture: all sites
Barad 2005 <sup>134</sup>	Cohort	OCs <sup>135</sup>		1.07 (1.01–1.15)	4,674	80,947	First fracture
Barad 2005 <sup>136</sup>	Cohort	OCs		1.15 (1.04-1.27)	4,674	80,947	First fracture
Barad 2005 <sup>137</sup>	Cohort	OCs		1.09 (0.97–1.23)	4,674	80,947	First fracture
Lanza 2013 <sup>138</sup>	Retrospective cohort study	DMPA <sup>139</sup>		1.41 (1.35–1.47)	11,822	312,395	Incident fractures

<sup>127</sup> From the Royal College of General Practitioners (RCGP) Oral Contraception Study.

<sup>128</sup> OC use > 97 months vs no use. Recruited age 25 to 39 years; followed to 45 years.

<sup>129</sup> OC use > 97 months vs no use. Recruited age 25 to 39 years; followed to 45 years.

<sup>130</sup> Interval since use: 73 to 96 months vs no use (radius or ulna). Recruited age 25 to 39 years; followed to 45 years.

<sup>131</sup> < 12 months vs no use (all fractures). Recruited age 25 to 39 years; followed to 45 years.

<sup>132</sup>  $\chi^2$  trend.

<sup>133</sup>  $\chi^2$  trend.

<sup>134</sup> Recruited age 50 to 74 years; OC use: any vs none.

<sup>135</sup> The patients were asked about oral contraceptive use, which likely was predominantly COCs but was not broken down with regard to COCs or POCs.

<sup>136</sup> Among women without any postmenopausal hormone treatment, past OC use for 5 years or less.

<sup>137</sup> Among women without any postmenopausal hormone treatment, past OC use for more than 5 years.

<sup>138</sup> They note that, “Although DMPA users experienced more fractures than nonusers, this association may be the result of confounding by a pre-existing higher risk for fractures in women who chose DMPA for contraception.” However, this is based on analysis of relatively few fractures prior to DMPA use.

<sup>139</sup> Depot medroxyprogesterone acetate = DMPA.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	Past use <sup>140</sup>	DMPA		1.32 (1.24–1.41)				Incident fractures
	Recent use <sup>141</sup>	DMPA		1.41 (1.31–1.50)				Incident fractures
	Current use <sup>142</sup>	DMPA		1.51 (1.41–1.61)				Incident fractures
Tuppurainen 1993 <sup>143</sup>	Case-control	OCs	1.21 (0.93–1.57)			629	13,100	All fractures
Tuppurainen 1993 <sup>144</sup>	Case-control	OCs	1.35 (0.88–2.05)			210	13,100	Wrist fractures
O’Neill 1996	Case-control	OCs	0.3 (0.1–0.9)			62	116	Distal forearm fractures only Population controls
O’Neill 1996	Case-control	OCs	0.7 (0.2–2.4)			62	50	Distal forearm fractures only Fall controls
Michaëlsson 1999 <sup>145</sup>	Case-control	Any <sup>146</sup>	0.75 (0.59–0.96)			1327	3312	Hip fractures
Vestergaard 2006 <sup>147</sup>	Case-control	OCs	<0.3 DDD/day	0.3–0.99 DDD/day	1+ DDD/day	64,548	193,641	Any fracture in the year 2000
	<25 years <sup>148</sup>	OCs	0.97 (0.91–1.03)	0.96 (0.92–1.01)	0.92 (0.86–0.98)			Any fracture in the year 2000
	25–49 years	OCs	0.91 (0.82–1.00)	0.90 (0.77–1.05)	0.87 (0.64–1.18)			Any fracture in the year 2000
	50+ years	OCs	0.92 (0.77–1.10)	0.69 (0.45–1.05)	0.62 (0.27–1.41)			Any fracture in the year 2000
Vestergaard 2008a <sup>149</sup>	Case-control	OCs	<0.3 DDD/day	0.3–0.99 DDD/day	1+ DDD/day	64,548	193,641	Any fracture in the year 2000
	<15	OCs	1.02 (0.75–1.37)	1.17 (1.01–1.37)	0.97 (0.85–1.11)			Any fracture in the year 2000

<sup>140</sup> Active DMPA use based on the interleaving of active 90-day exposures generated by each injection.

<sup>141</sup> Recent exposure is 640 or fewer days after the last active exposure.

<sup>142</sup> Past exposure begins after “recent” exposure (641 or more days after the last active exposure).

<sup>143</sup> Oral contraceptive use for 6+ years.

<sup>144</sup> Oral contraceptive use for 6+ years.

<sup>145</sup> No significant correlation was seen with duration of use, time since last use or time between last use and menopause.

<sup>146</sup> Any type of chemical contraceptive was evaluated, not separated as COCs or POCs.

<sup>147</sup> “The exposure time for oral contraceptives may thus maximally have spanned 5 years (from January 1, 1996, to December 31, 2000).” This and the other Vestergaard study are not useful as they do not take into account remote use or cumulative lifetime use. ORs shown.

<sup>148</sup> Defined daily dosages = DDD.

<sup>149</sup> Similar to Vestergaard 2006; only looked at use within the past 5 years. A younger group examined here. ORs shown.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	15.1-17	OCs	1.22 (1.02–1.47)	1.14 (1.00–1.30)	1.04 (0.90–1.19)			Any fracture in the year 2000
	17.1-19	OCs	0.97 (0.87–1.09)	0.93 (0.84–1.02)	1.02 (0.89–1.18)			Any fracture in the year 2000
	>19	OCs	0.99 (0.93–1.05)	1.00 (0.93–1.08)	0.88 (0.78–0.99)			Any fracture in the year 2000
Vestergaard 2008b <sup>150</sup>	Case-control	DMPA	1.44 (1.01–2.06)			64,548	193,641	Any fracture in the year 2000 DMPA use
Wei 2011 <sup>151</sup>	Cross-sectional		<5 years of use	5-10 years of use	>10 years of use		491	
		OCs	0.85 (0.45–1.58)	0.45 (0.21–0.93)	0.75 (0.36–1.54)			Presence of vertebral deformity
		OCs	0.96 (0.62–1.48)	0.63 (0.37–1.07)	0.94 (0.56–1.56)			Number of vertebral deformities
Meier 2010 <sup>152</sup>	Case-control		Current Use	Past Use		17,527	70,130	Incident fracture
	1-2 DMPA Scripts	DMPA	1.18 (0.93–1.49)	1.17 (1.07–1.29)				Incident fracture
	3-9 DMPA scripts	DMPA	1.36 (1.15–1.60)	1.23 (1.11–1.36)				Incident fracture
	10+ DMPA scripts	DMPA	1.54 (1.33–1.78)	1.30 (1.09–1.55)				Incident fracture
	1-2 COC Scripts	COCs	1.01 (0.87–1.18)	1.00 (0.95–1.07)				Incident fracture
	3-9 COC scripts	COCs	1.01 (0.94–1.09)	0.99 (0.94–1.04)				Incident fracture
	10+ COC scripts	COCs	1.09 (1.03–1.16)	1.03 (0.97–1.10)				Incident fracture
Memon 2011 <sup>153</sup>	Case-control	COCs	1.05 (0.86–1.29)			651	1302	Any fracture
Kyvernitakis 2016 <sup>154</sup>	Case-control		OR Current Use	OR Past Use		4189	4189	First-time fracture diagnosis
	1-2 DMPA scripts	DMPA	0.97 (0.51–1.86)	0.96 (0.73–1.26)				
	3-9 DMPA scripts	DMPA	2.41 (1.42–4.08)	1.14 (0.86–1.51)				
	10+ DMPA scripts	DMPA	1.46 (0.96–2.23)	1.55 (1.07–2.27)				

<sup>150</sup> Similar to Vestergaard 2006; only looked at use within the past 5 years. DMPA examined here. ORs shown.

<sup>151</sup> Small cross-sectional study. ORs shown.

<sup>152</sup> Females aged 20–44 years with an incident fracture diagnosis between 1995 and 2008.

<sup>153</sup> Nested case-control study of the Cooper study from the Royal College of General Practitioners (RCGP) Oral Contraception Study. Last OC use > 10 years vs never.

<sup>154</sup> Women between 20 and 44 years of age with a first-time fracture diagnosis, matched with random controls using the Disease Analyzer database.

Study	Study Design	Intervention	OR	RR		Cases	Controls or Cohort Size	Outcome
	1-2 COC scripts	COCs	0.98 (0.73–1.31)	0.90 (0.77–1.05)				
	3-9 COC scripts	COCs	1.39 (1.12–1.73)	0.90 (0.78–1.03)				
	10+ COC scripts	COCs	1.07 (0.88–1.30)	1.04 (0.90–1.21)				

## Osteoporotic Fracture References

- Barad D, Kooperberg C, Wactawski-Wende J, Liu J, Hendrix SL, and Watts NB. Prior oral contraception and postmenopausal fracture: a Womens' Health Initiative observational cohort study. *Fertility and Sterility* 2005; 84:374–383.
- Cooper C, Hannaford P, Croft P, and Kay CR. Oral contraceptive pill use and fractures in women: a prospective study. *Bone* 1993; 14(1):41–45.
- Curtis KM and Martins SL. Progestin-only contraception and bone mineral density: a systematic review. *Contraception* 2006; 73:470–487.
- FDA 2004: U.S. Food and Drug Administration. FDA Talk Paper. Black box Warning Added Concerning Long-term Use of Depo-Provera Contraceptive Injection. <http://web.archive.org/web/20070809090332/http://www.fda.gov/bbs/topics/ANSWERS/2004/ANS01325.html> (accessed 2019 February 23).
- Lanza LL, McQuay LJ, Rothman KJ, Bone HG, Kaunitz AM, Harel Z, Ataher Q, Ross D, Arena PL, and Wolter KD. Use of depot medroxyprogesterone acetate contraception and incidence of bone fracture. *Obstetrics and Gynecology* 2013; 121(3):593–600.
- Lopez LM, Chen M, Mullins Long S, Curtis KM, and Helmerhorst FM. Steroidal contraceptives and bone fractures in women: evidence from observational studies (Review). *Cochrane Database of Systematic Reviews* 2015; Issue 7. Art. No.: CD009849. DOI: 10.1002/14651858.CD009849.pub3.
- Meier C, Brauchli YB, Jick SS, Kraenzlin ME, and Meier CR. Use of depot medroxyprogesterone acetate and fracture risk. *Journal of Clinical Endocrinology and Metabolism* 2010; 95(11):4909–4916.
- Memon S, Iversen L, and Hannaford PC. Is the oral contraceptive pill associated with fracture in later life? New Evidence from the Royal College of General Practitioners Oral Contraception Study. *Contraception* 2011; 84(1):40–47.
- Michaelsson K, Baron JA, Farahmand BY, and Ljunghall S. Influence of parity and lactation on hip fracture risk. *American Journal of Epidemiology* 2001; 153(12):1166–1172.
- Michaelsson K, Baron JA, Farahmand BY, Persson I, and Ljunghall S. Oral-contraceptive use and risk of hip fracture: a case-control study. *Lancet* 1999; 353(9163):1481–1484.
- O'Neill TW, Marsden D, Adams JE, and Silman AJ. Risk factors, falls, and fracture of the distal forearm in Manchester, UK. *Journal of Epidemiology and Community Health* 1996; 50(3):288–292.
- Parazzini G, Tavani A, Ricci E, and La Vecchia C. Menstrual and reproductive factors and hip fractures in post menopausal women. *Maturitas* 1996; 24:191–196.
- Sirola J, Rikkonen T, Tuppurainen M, Honkanen R, and Kroger H. Should risk of bone fragility restrict weight control for other health reasons in postmenopausal women?—A ten year prospective study. *Maturitas* 2012; 71(2):162–168.

Tuppurainen M, Honkanen R, Kröger H, Saarikoski S, and Alhava E. Osteoporosis risk factors, gynaecological history and fractures in perimenopausal women—the results of the baseline postal enquiry of the Kuopio Osteoporosis Risk Factor and Prevention Study. *Maturitas* 1993; 17(2):89–100.

Vessey M, Mant J, and Painter R. Oral contraception and other factors in relation to hospital referral for fracture. Findings in a large cohort study. *Contraception* 1998; 57(4):231–235.

Vestergaard P, Rejnmark L, and Mosekilde L. Fracture risk in very young women using combined oral contraceptives. *Contraception* 2008a; 78(5):358–364.

Vestergaard P, Rejnmark L, and Mosekilde L. Oral contraceptive use and risk of fractures. *Contraception* 2006; 73(6):571–576.

Vestergaard P, Rejnmark L, and Mosekilde L. The effects of depot medroxyprogesterone acetate and intrauterine device use on fracture risk in Danish women. *Contraception* 2008b; 78(6):459–464.

Wei S, Venn A, Ding C, Foley S, Laslett L, and Jones G. The association between oral contraceptive use, bone mineral density and fractures in women aged 50-80 years. *Contraception* 2011; 84(4):357–362.

Women's Health Initiative Study Group. Design of the Women's Health Initiative clinical trial and observational study. *Controlled Clinical Trials* 1998; 19(1):61–109.

## Impact of Contraceptives on Body Mass

Weight gain is a common complaint among contraceptive users but whether use of contraceptives is causally related remains undefined. Progestin-only contraceptives are most commonly associated with weight gain complaints and discontinuation. A recent Cochrane review (Gallo et al. 2014) examined the effect of combined oral contraceptives on weight gain and concluded existing data does not support a causal relationship. A second review of progestin-only contraceptives on weight gain (Lopez et al. 2016) found most studies of low to moderate quality but did conclude weight gain of up to 2kg (4.4 lbs) within the first year of use with continued increases thereafter. The authors advised appropriate counselling on expected weight changes to minimize discontinuation due to perceived weight gain.

The attached table (Table 13) summarizes studies of 1 year or longer that examined weight and body mass changes in contraceptive users in comparison to non-hormonal contraceptives or no method. Several additional studies compare various contraceptives for their effect on weight or body composition, but these do not directly address our focus.

The strongest data appear to be the deleterious effects of levonorgestrel-releasing IUDs on percent lean and fat body mass. Total body weight change does not appear different between groups and several large studies have shown no significant differences. However, a significant increase in % fat mass with a corresponding decrease in % lean body mass was observed in both studies where these were measured. A similar effect was seen from oral desogestrel in a single study.

Thus, while limited to date, data suggest that use of progestin-only contraceptives may have deleterious effects on % fat and % lean body mass with no significant overall effect on total body weight.

A review of current Mirena labeling makes no mention of changes in lean or fat body mass composition.

Retrospective, but not more recent, prospective studies also show DMPA use is associated with significant gains in weight. The data appear too mixed to draw firm conclusions.



Table 13 – Effect of Chemical Contraceptives on Weight Gain

Study	Design	Comparison	N	Time	Weight change (Kg)	Fat mass change	Lean mass change	Comments
Pantoja 2010	Retrospec.	DMPA 150 vs CuIUC	758	1yr	1.76 vs-0.42*			Largest differences noted in normal and overweight BMI subgroups, minimal differences in obese BMI subgroup
				2yr	3.1 vs 0.4*			
				3yr	3.9 vs 0.8*			
Modesto 2015	Retrospec.	DMPA150 vs CuIUC	1277	1yr	1.3 vs 0.2*			Adjusted for years of school & # children. 20% loss @4yrs 84% @ 10yr.
				4yr	3.5 vs 1.9*			
				10yr	6.6 vs 4.9*			
Taneepanichskul 1998	Retrospec.	DMPA 150 vs CuIUC	100	10yr	10.9 vs 11.2			Included women 37-50 years (no younger women)
Vickery 2013	Prospec.	DMPA 150 vs CuIUC	167	1yr	2.2 vs 0.16			CHOICE study subgroup
Dal'Ava 2014	Prospec.	DMPA 150 vs CuIUC	110	1yr	1.9vs 1.1	1.6 vs -0.9 (Kg)	0.3 vs 1.2 (kg)	Paired by age (+/-2yr) & weight (+/-2kg)
Dos Santos 2014	Prospec.	DMPA 150 vs CuIUC	71	1yr	1.4 vs 0.3	1.57 vs 0.52 (kg)	(0.31) vs (0.26) (kg)	Matched by age & BMI ( )= negative value
<b>Studies comparing LNG IUC to non-hormonal contraceptive</b>								
Study	Design	Comparison	N	Time	Weight change (Kg)	Total body fat	Lean body mass	
Dal'Ava 2012	Prospec.	LNG-IUC vs non-hormonal IUC	76	1yr	2.9 vs 1.4	2.5% vs -1.3%*	(1.4%) vs 1.0%*	Paired by age & BMI
Napolitano 2015	Prospec.	LNG IUC vs no method	60	1yr	0.6 vs (0.2)	1.1% vs (0.5%)*	(1.1%) vs 0.5*	
Vickery 2013	Prospec.	LNG-IUC vs Cu IUC	230	1yr	1.03 vs 0.16	nd	nd	

Modesto 2015	Retrospec.	LNG-IUC vs CuIUC	1204	1yr	0.7 vs 0.2	nd	nd	
				4yr	2.7 vs 1.9			
				10yr	4.0 vs 4.9			

**Studies comparing progestin-only COCs to non-hormonal**

Study	Design	Comparison	N	Time	Weight change (Kg)	Total body fat	Lean body mass	
Napolitano 2015	Prospec.	Desogestrel 75ug vs no hormonal	68	1yr	0.3 vs -0.2	1.1% vs -0.5%*	(2.8%) vs 0.5%*	

**Studies comparing combined COCs to non-hormonal**

None found-

Abstract from 2014 Cochrane review of combined oral contraceptives on weight gain:

"We found 49 trials that met our inclusion criteria. The trials included 85 weight change comparisons for 52 distinct contraceptive pairs (or placebos). *The four trials with a placebo or no intervention group did not find evidence supporting a causal association* between combination oral contraceptives or a combination skin patch and weight change. Most comparisons of different combination contraceptives showed no substantial difference in weight. In addition, discontinuation of combination contraceptives because of weight change did not differ between groups where this was studied.

Gallo MF, Lopez LM, Grimes DA, Carayon F, Schulz KF, Helmerhorst FM. Combination contraceptives: effects on weight. Cochrane Database of Systematic Reviews 2014, Issue 1. Art. No.: CD003987.

\* Significant difference (p<0.05).

## Impact of Contraceptives on Body Mass References

Dal'Ava N, Bahamondes L, Bahamondes MV, Bottura BF, and Monteiro I. Body weight and body composition of depot medroxyprogesterone acetate users. *Contraception* 2014; 90(2):182–187.

Dal'Ava N, Bahamondes L, Bahamondes MV, de Oliveira Santos A, and Monteiro I. Body weight and composition in users of levonorgestrel-releasing intrauterine system. *Contraception* 2012; 86(4):350–353.

Dos Santos, PdeNS, Modesto WO, Dal'Ava N, Bahamondes M, Pavin E, and Fernandes A. Body composition and weight gain in new users of the three-monthly injectable contraceptive, depot-medroxyprogesterone acetate, after 12 months of follow-up. *European Journal of Contraception and Reproductive Health Care* 2014; 19(6):432–438.

Gallo MF, Lopez LM, Grimes DA, Carayon F, Schulz KF, and Helmerhorst FM. Combination contraceptives: effects on weight. *Cochrane Database of Systematic Reviews* 2014; Issue 1. Art. No.: CD003987.

Lopez LM, Ramesh S, Chen M, Edelman A, Otterness C, Trussell J, and Helmerhorst FM. Progestin-only contraceptives: effects on weight (Review). *Cochrane Database of Systematic Reviews* 2016; Issue 8. Art. No.: CD008815.

Lopez LM, Ramesh S, Chen M, Edelman A, Otterness C, Trussell J, and Helmerhorst FM. Progestin-only contraceptives: effects on weight. *Cochrane Database Syst Reviews* 2016a Aug 28;(8):CD008815. Doi: 10.1002/14651858.

Modesto W, dos Santos PdeNS, Correia VM, Borges L, and Bahamondes L. Weight variation in users of depot-medroxyprogesterone acetate, the levonorgestrel-releasing intrauterine system and a copper intrauterine device for up to ten years of use. *European Journal of Contraception and Reproductive Health Care* 2015; 20(1):57–63.

Napolitano A, Zanin R, Palma F, Romani C, Grandi G, Di Carlo C, and Cagnacci A. Body composition and resting metabolic rate of perimenopausal women using continuous progestogen contraception. *European Journal of Contraception and Reproductive Health Care* 2015 Aug 25 [Epub ahead of print]:1–8.

Pantoja M, Medeiros T, Baccarin MC, Morais SS, Bahamondes L, and Fernandes AM. Variations in body mass index of users of depot-medroxyprogesterone acetate as a contraceptive. *Contraception* 2010; 81(2):107–111. Taneepanichskul S, Reinprayoon D, Khaosaad P. Comparative study of weight change between long-term DMPA and IUD acceptors. *Contraception* 1998; 58(3):149–151.

Vickery Z, Madden T, Zhao Q, Secura GM, Allsworth JE, and Peipert JF. Weight change at 12 months in users of three progestin-only contraceptive methods. *Contraception* 2013; 88(4):503–508.

## Urogenital Effects of Contraceptives

In addition to cervical cancer and interstitial cystitis, noted above, there are other adverse urogenital effects of COCs that should be communicated to patients. These include bacteriuria (Zahran 1976; calculated OR 3.57), urinary tract infection (Engel 1979: 27–50% incidence), bladder trabeculation (Zahran 1976; calculated OR 11.7), recurrent vulvovaginal candidiasis (Spinillo 1995, Yusuf 2007; OR 2.08), vaginal dryness (Lee 2017), vulvar vestibulitis (Champaneria 2016: OR 2.1 95 % CI 1.26–3.49; also noted in Lee 2017), and Female Sexual Dysfunction (FSD) (Lee 2017). FSD appears related to OC-induced dyspareunia, reduced sexual desire and libido (Lee 2017). This risk is increased if COCs are used in adolescents and the duration of OC use is at least 2 years (Lee 2017), although some newer COCs containing drospirenone 3 mg plus EE 30 mg and gestodene 75 mg plus EE 20 mg appear to have a reduction in these risks (Lee 2017).

These urogenital risks, especially FSD where there is substantial literature, should be referenced in prescribing information and patient pamphlets.

## Urogenital Effects References

Champaneria R, D'Andrea RM, and Latthe PM. Hormonal contraception and pelvic floor dysfunction: a systematic review. *Int Urogynecol J*. 2016; 27:709–722.

Engel HJ. Adverse effects of oral contraceptives. *Med Monatsschr Pharm*. 1979:199–204.

Lee JJML, Low LL, and Ang SB. Oral Contraception and Female Sexual Dysfunction in Reproductive Women. *Sexual Medicine Reviews*. 2017; 5:31–44.

Spinillo A, Capuzzo E, Nicola S, and Baltaro F, Ferrari A, Monaco A. The impact of oral contraception on vulvovaginal candidiasis. *Contraception*. 1995; 51:293–297.

Yusuf A, Chowdhury AQ, Sattar ANI, and Rahman M. Evaluation of the Effect of Contraceptives in Prevalence of Candida Species on Vaginal Candidiasis in Dhakam Bangladesh. *Bangladesh J Med Microbiol*. 2007; 01:61–64.

Zahran MM, Osman MI, Kamel M, Fayad M, Mooro H, and Youssef AF. Effects of contraceptive pills and intrauterine devices on urinary bladder. *Urology*. 1976; 8:567–574.

## Venous Thromboembolism and Contraceptives

The current language on the black box warning of certain contraceptives regarding risk of cardiovascular events clearly misleads women about the real risks of these drugs. It says: WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS. A study (Gomer 2009) conducted among 300 women concluded “that most of them believe that certain risks are only associated with being over 35 years of age and/or smoking.” Instead, the label should clearly state that anyone taking the medications without good knowledge of the risk factors could experience a potentially life-threatening cardiovascular event and should discuss the risks with a medical provider.

The incidence of venous thromboembolism (VTE) for healthy women can significantly increase with the use of hormonal contraceptives, even women under 35 and not-smoking. In a 2012 article about birth control side effects, Dr. Rebecca Peck (Peck R 2012) reports that “Oral contraceptives are associated with a three to five times higher risk of VTE (Van Hylckama VA 2009).” Third and fourth generation combined hormonal contraceptives (CHC) have been found to put women at an even much higher risk, leading to major lawsuits against some manufacturers and changes in regulations in several countries. In his opinion published in Drug Safety, Dr. Lidegaard, the author of several studies on the subject, states: “Of 14 studies specifically assessing the risk in users of CHC with desogestrel or gestodene, 13 found a higher risk with use of these products when compared to the use of CHC with levonorgestrel” (Lidegaard 2014). Drospirenone, the progestin contained in Yaz and Jasmine, also increases the risk of VTE over levonorgestrel by a factor of 1.5 to 2.8. “The relative risk [of Drospirenone was 6.3 as compared with nonusers in both the large Dutch (Van Hylckama 2009) and Danish (Lidegaard 2011) study.” The author comments that “the studies demonstrating risk differences between CHC with different progestins are generally methodologically more transparent and more robust than those demonstrating no difference, especially concerning exclusion of women with predispositions for VTE.” Another large study published in 2015 (Vinogradova 2015) reviewed 10,552 cases of VTE reported between 2001 and 2013 in the UK and found similar elevated risks of VTE with these CHC: “Corresponding risks associated with current exposure to desogestrel (4.28, 3.66 to 5.01), gestodene (3.64, 3.00 to 4.43), drospirenone (4.12, 3.43 to 4.96), and cyproterone (4.27, 3.57 to 5.11) were significantly higher than those for second generation contraceptives levonorgestrel (2.38, 2.18 to 2.59).” Note that the odds ratios were “adjusted for smoking status, alcohol consumption, ethnic group, body mass index, comorbidities, and other contraceptive drugs.”

Most importantly, the risk levels are multiplied if women have other risk factors. For instance, women who have the genetic blood condition known as Factor V Leiden could have a risk as high as 18 per 10,000 woman-years. If these women stay on the product for 10 years, their risks could be 250 per 10,000 woman-years, or 2.5% as risks increase with aging (Lidegaard 2014).

Dr. Lidegaard concludes: “Therefore, women with known risk factors of VTE are advised to be reluctant to use CHC. The relative risk of VTE with different dispositions is as follows: previous thrombosis: > 50 (Le Moigne 2013), genetic abnormalities such as factor V Leiden mutation (heterozygous): 6, deficiency of protein C: 10, of protein S: 10, of antithrombin: 25, and of prothrombin 20210A: 3 (Phillippe 2014). Pregnancy with delivery on average: 8, adiposity: 2–3 and immobilization 2–5 depending on how long time you are immobilized. Family disposition (first-degree relatives with VTE before their 50th year) doubles the risk of VTE. Women with such dispositions are generally recommended to use progestin-only contraception, which does not increase the risk of VTE except perhaps for medroxyprogesterone depots. A genetic screening should until further also be restricted to women with a family disposition” (Lidegaard 2014).

In a 2018 systematic review (Keenan 2018) of the most evidenced-based articles from the 1960s to 2018 comparing users of COCs to nonusers, with a confirmed diagnosis of VTE, and including more than 17 million

woman-years of observation, women on HC increase their risk by 3- to 9-fold. However, the first year of use has the highest risk for clot formation, and if a woman is younger than 30, her risk is increased 13-fold in the first year. Obesity can increase the risk of being on hormonal contraception, about doubling the risk compared to a woman of normal weight on the pill. It is not considered cost-effective to check for thrombophilia, a genetic disposition to form blood clots, but for those with thrombophilia, the risk can be as high as 62-fold in the first year.

This systematic review of the literature concludes that 136–260 women die from VTE a year in the United States from hormonal contraception. Combined with the added risk of stroke and heart attack from the COCs, 300–400 women die each year in the United States simply due to their choice of using HC for family planning (Keenan 2018). To give some perspective, meningitis killed 45 people (of all ages) in 2017: most US States mandate meningitis vaccination for college and university students.

A summary of studies is shown in Table 14.

Table 14 – Relative Risk of Venous Thromboembolism in Current Users of Different Combined Hormonal Contraceptives as Compared with Nonusers Unless Otherwise Specified

Study	Data Sampling Period	VTE (number)	CHCs with levonorgestrel RR (95% CI)	CHCs with desogestrel/gestodene RR (95% CI)	CHCs with drospirenone RR (95% CI)
Blomenkamp 1995	1988 - 1992	126	3.8 (1.7 - 8.4)	8.7 (3.9 - 19.3)	-
WHO 1995a, 1995b	1989 - 1993	433	3.6 (2.5 - 5.1)	7.4 (4.2 - 12.9)	-
Jick 1995	1991 - 1994	80	1 (reference)	1.8 (1.0 - 3.2)	-
Spitzer 1996	1991 - 1995	471	3.7 (2.2 - 6.2)	6.7 (3.4 - 13.0)	-
Lewis 1999	1993 - 1995	502	2.9 (1.9 - 4.2)	2.3 (1.5 - 3.5)	-
Farmer 1997	1991 - 1995	85	3.1‡ (2.1 - 4.5)	5.0‡ (3.7 - 6.5)	-
Todd 1999	1992 - 1997	99	1 (reference)	1.4 (0.7 - 2.8)	-
Bloemenkamp 1999	1994 - 1998	185	3.7 (1.9 - 7.2)	5.6 (not given)	-
Parkin 2000	1990 - 1998	26	5.1 (1.2 - 21.4)	14.9 (3.5 - 64.3)	-
Lidegaard 2002	1994 - 1998	987	2.9 (2.2 - 3.8)	4.0 (3.2 - 4.9)	-
Dinger 2007	2000 - 2004	118	1 (reference)	1.3 (NA)	1.0 (0.6 - 1.8)
Vlieg 2009	1999 - 2004	1524	3.6 (2.9 - 4.6)	7.3 (5.3 - 10.0)/5.6 (3.7 - 8.4)	6.3 (2.9 - 13.7)
Lidegaard 2009	1995 - 2005	4213	2.0 (1.8 - 2.3)	3.6 (3.3 - 3.8)	4.0 (3.3 - 4.9)
Dinger 2010	2002 - 2008	680	1 (reference)	NA	1.0 (0.6 - 1.8)
Parkin 2011	2002 - 2009	61	1 (reference)	NA	2.7 (1.5 - 4.7)
Jick 2011	2002 - 2008	186	1 (reference)	NA	2.8 (2.1 - 3.8)
Lidegaard 2011	2001 - 2009	4246	2.2 (1.7 - 2.8)	4.2 (3.6 - 4.9)	4.5 (3.9 - 5.1)
Confirmed only	2001 - 2009	2707	2.9 (2.2 - 3.8)	6.8 (5.7 - 8.1)	6.3 (5.4 - 7.5)
FDA Kaiser 2011	2001 - 2007	625	1 (reference)	NA	1.5 (1.2 - 1.9)
Gronich 2011	2002 - 2008	518	1 (reference)	1.4 (0.9 - 2.1)	1.7 (1.0 - 2.7)
Lidegaard 2012	2001 - 2010	5287	3.2 (2.7 - 3.8)	6.5 (4.7 - 8.9)*	NA
Dinger 2014	2005 - 2010	162	1 (reference)	NA	0.8 (0.5 - 1.6)

‡ Absolute risk per 10,000 years.

\* Vaginal ring with the third-generation progestin etonogestrel.

## Venous Thromboembolism References

Bloemenkamp KWM, Rosendaal FR, Büller HR, Helmerhorst FM, Colly LP, and Vandenbroucke JP. Risk of venous thrombosis with use of current low-dose oral contraceptives is not explained by diagnostic suspicion and referral bias. *Arch Intern Med* 1999; 159:65–70.

Bloemenkamp KWM, Rosendaal FR, Helmerhorst FM, Büller HR, and Vandenbroucke JP. Enhancement by factor V Leiden mutation of risk of deep-vein thrombosis associated with oral contraceptives containing a third-generation progestagen. *Lancet* 1995; 346:1593–1596.

Dinger J, Assmann A, M€ohner S, and Minh TD. Risk of venous thromboembolism and the use of dienogest- and drospirenone-containing oral contraceptives: results from a German case-control study. *J Fam Plann Reprod Health Care* 2010; 36:123–129.

Dinger J, Bardenheuer K, and Heinemann K. Cardiovascular and general safety of a 24-day regimen of drospirenone-containing combined oral contraceptives: final results from the International Active surveillance Study of Women Taking Oral Contraceptives. *Contraception* 2014; 89(4):253–263.

Dinger JC, Heinemann LAJ, and Kuhl-Habich D. The safety of a drospirenone-containing oral contraceptive: final results from the European Active Surveillance study on oral contraceptives based on 142,475 women years of observation. *Contraception* 2007; 75:344–354.

Farmer RDT, Lawrenson RA, Thompson CR, Kennedy JG, and Hambleton IR. Population-based study of risk of venous thromboembolism associated with various oral contraceptives. *Lancet* 1997; 349:83–88.

Food and Drug Administration, Office of surveillance and epidemiology. Combined hormonal contraceptives (CHCs) and the risk of cardiovascular disease endpoints. FDA. 2011; <http://www.fda.gov/downloads/Drugs/DrugSafety/UCM277384.pdf>.

Gomer K. Women, Birth Control Pills, and Thrombophilia: An Analysis of Risk Communication Kerry Gomer, Clemson University. 2009; [https://tigerprints.clemson.edu/cgi/viewcontent.cgi?referer=&httpsredir=1&article=1573&context=all\\_theses](https://tigerprints.clemson.edu/cgi/viewcontent.cgi?referer=&httpsredir=1&article=1573&context=all_theses).

Gronich N, Lavi I, and Rennert G. Higher risk of venous thrombosis associated with drospirenone-containing oral contraceptives: a population-based cohort study. *CMAJ* 2011; 183(18):E1319-25.

Jick H, Jick SS, Gurewich V, Myers MW, and Vasilakis C. Risk of idiopathic cardiovascular death and nonfatal venous thromboembolism in women using oral contraceptives with differing progestagen components. *Lancet* 1995; 346:1589-1593.

Jick SS, and Hernandez RK. Risk of non-fatal venous thromboembolism in women using oral contraceptives containing drospirenone compared with women using oral contraceptives containing levonorgestrel: case-control study using United States claims data. *British Medical Journal* 2011; 340:d2151.

Keenan L, Kerr T, Duane M, and Van Gundy K. Systematic Review of Hormonal Contraception and Risk of Venous Thrombosis. *The Linacre Quarterly* 2018; 85(4):470–477.



- Le Moigne E, Delluc A, Tromeur C, Nowak E, Mottier D, Lacut K, and Le Gal G. Risk of recurrent venous thromboembolism among young women after a first event while exposed to combined oral contraception versus not exposed to: a cohort study. *Thromb Res* 2013; 132:51–55.
- Lewis MA, MacRae KD, Kúhl-Habich D, Bruppacher R, Heinemann LA, and Spitzer WO. The differential risk of oral contraceptives: the impact of full exposure history. *Hum Re prod* 1999; 14:1493–1499.
- Lidegaard Ø, Edström B, and Kreiner S. Oral contraceptives and venous thromboembolism. A five-year national case-control study. *Contraception* 2002; 65:187–196.
- Lidegaard Ø, Løkkegaard E, Svendsen AL, and Agger C. Hormonal contraception and risk of venous thromboembolism: national follow-up study. *British Medical Journal* 2009; 339:b2890.
- Lidegaard Ø, Nielsen LH, Skovlund CW, and Løkkegaard E. Venous thrombosis in users of non-oral hormonal contraception: follow-up study, Denmark 2001–10. *British Medical Journal* 2012; 344:e2990.
- Lidegaard Ø, Nielsen LH, Skovlund CW, Skjeldestad FE, and Løkkegaard E. Risk of venous thromboembolism from use of oral contraceptives containing different progestogens and estrogen doses: Danish cohort study 2001–2009. *British Medical Journal* 2011; 343:d6423.
- Lidegaard Ø. Hormonal contraception, thrombosis and age. *Expert Opin Drug Saf* 2014; 13:1353–360.
- Parkin L, Sharples K, Hernandez RK, and Jick SS. Risk of venous thromboembolism in users of oral contraceptives containing drospirenone or levonorgestrel: nested case-control study based on UK General Practice Research Database. *British Medical Journal* 2011; 340:d2139.
- Parkin L, Skegg DCG, Wilson M, Herbison GP, and Paul C. Oral contraceptives and fatal pulmonary embolism. *Lancet* 2000; 355:2133–2134.
- Peck R and Norris CW. Significant Risks of Oral Contraceptives (OCs) *Linacre Q* 2012; 79(1): 41–56. Published online 2012 Feb 1. Doi: 10.1179/002436312803571447.
- Phillippe HM, Hornsby LB, Treadway S, Armstrong EM, and Bellone JM. Inherited thrombophilia. *J Pharm Pract* 2014; 27:227–233.
- Spitzer WO, Lewis MA, Heinemann LAJ, Thorogood M, and MacRae KD. Third generation oral contraceptives and risk of venous thromboembolic disorders: an international case-control study. *British Medical Journal* 1996; 312:83-88.
- Todd J-C, Lawrenson R, Farmer RDT, Williams TJ, and Leydon GM. Venous thromboembolic disease and combined oral contraceptives: a re-analysis of the MediPlus database. *Hum Reprod* 1999; 14:1500–1505.
- Van Hylckama VA, Helmerhorst FM, Vandenbroucke JP, Doggen CJ, and Rosendaal FR. The Venous Thrombotic Risk of Oral Contraceptives, Effects of Estrogen Dose and Progestogen Type: Results of the MEGA Case-Control Study. *British Medical Journal* 2009; 339:b2921. Doi:10.1136/bmj.b2921.

Vinogradova Y, Coupland C and Hippisley-Cox J. Use of combined oral contraceptives and risk of venous thromboembolism: nested case-control studies using the QResearch and CPRD databases. *British Medical Journal* 2015; 350:h2135. Doi: <https://doi.org/10.1136/bmj.h2135> (Published 26 May 2015).

World Health Organisation Collaborative Study on Cardiovascular Disease and Steroid Hormone Contraception. Effect of different progestogens in low estrogen oral contraceptives on venous thromboembolic disease. *Lancet* 1995; 346:1582–1588.

World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Venous thromboembolic disease and combined oral contraceptives: results of international multicentre case-control study. *Lancet* 1995; 346:1575-1582.

Yasmin Prescribing Information: Highlights of Yasmin Prescription Information, [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2012/021098s019lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/021098s019lbl.pdf).

## Atherosclerosis and Cardiovascular Events

Noting that previous studies had demonstrated women on oral contraceptives (OC) faced a fourfold increased risk of heart attack (Hennekens 1977; Vessey 1976; Beral 1976), researchers in 1982 set out to understand the pathogenesis of vascular disease related to COCs. They found that combination oral contraceptives (COC) caused “greater cell proliferation and incorporation...in both human arterial smooth muscle cells and dermal fibroblasts.” Smooth muscle cell proliferation is an integral feature of all atherosclerotic lesions (Bagdade 1982).

In 2007, a presentation at the American Heart Association meeting described a study of 1,301 Belgian women, which showed that women had a 20 to 30 percent increase of plaque for every decade on COCs (Zoler 2007). They noted that active OC users had elevated C-reactive protein levels, three times higher than non-users. C-reactive protein is a biomarker for many inflammation-related arterial (and autoimmune) diseases, which was recently the subject of another presentation (Rietzschel 2018).

They evaluated the carotid and femoral pulse wave velocity (PWV) and found the average PWV among non-users was 6.6 m/sec, while the average among current OC users was 6.75 m/sec. The blood pressure of current OC users was also significantly higher (4.3/2.3 mm Hg higher than non-users) (Zoler 2007). Lead investigator Dr. Ernst Rietzschel said this study “changes our thinking about oral contraceptives just causing an increased thrombotic risk. Instead, it appears as though OC use may also cause long-term structural damage to the vasculature.” These findings were supported by an evaluation of large artery stiffness in the ENIGMA study (Hickson 2011) although other smaller studies have shown conflicting data (Yu 2014, Priest 2018).

A study of homocysteine and nitric oxide levels compared 50 healthy women with normal menstrual cycles as a control group and 50 healthy women receiving oral contraceptive pills for at least three menstrual cycles (Fallah 2012). They noted that after 3 months of treatment, homocysteine levels were significantly increased ( $P = 0.027$ ), and there was a significant and considerable decrease ( $P = 0.048$ ) in NO concentration of oral contraceptive pill (OCP) consumers. Another study evaluated the effect of COCs on homocysteine and C-reactive protein levels in women (Norouzi 2011). This observational cross-sectional analysis included 90 healthy, non-obese women (mean age 25 years). Forty-five healthy women on OCP and 45 healthy controls were studied. COC users had a minimum of 3 cycles on COCs. The results showed that the homocysteine ( $13.268 \pm 3.475$  vs.  $7.288 \pm 2.621$   $\mu\text{mol/L}$ ) and CRP ( $5863.0 \pm 1349.5$  vs.  $1138.3 \pm 691.12$  ng/ml) levels were significantly higher in women receiving OCP in comparison with the control group ( $p=0.027$  and  $p<0.001$ , respectively). Similarly, a cross-sectional study, in 2011-2012, evaluated 60 healthy premenopausal women (30 cases of COC consumers and 30 controls as nonconsumers), aged between 25 and 45 years who were current users for at least a 3-year period. They evaluated brachial artery endothelial function (using flow-mediated dilatation (FMD)) and common carotid artery intima-media thickness (Heidarzadeh 2014). They noted that there was a significant FMD% difference between 2 groups of cases and controls:  $11 \pm 3.53$  versus  $15.80 \pm 9.22$  ( $P = 0.01$ ). In addition, a significant mean CCA-IMT thickness difference was detected:  $0.53 \pm 0.07$  versus  $0.44 \pm 0.08$  ( $P = 0.00$ ). Although these results were not significant after multiple regression analysis, the authors noted that their results were in favor of early atherosclerotic changes in prolonged users of COCs.

The Danish Heart Association released the results of a 15-year historic cohort study looking at thrombotic stroke and myocardial infarction, which observed over 1.6 million women. The results demonstrated that women taking COCs with ethinyl estradiol at a dose of 20  $\mu\text{g}$  had a risk of arterial thrombosis that was 0.9 to 1.7 times higher than non-users, while those taking a dose of 30 to 40  $\mu\text{g}$  had a 1.3 to 2.3 higher risk (Lidegaard 2012). The risk of thrombotic stroke appeared to be independent of duration of use, while the risk for myocardial infarction increased with duration of use (Table 15).

Together, these studies suggest that protracted use of COCs can induce atherosclerotic changes independent of any pro-thrombotic effect. These changes may contribute to the increase in thrombotic stroke and myocardial infarction seen in COC users.

*Table 15 – Relative Risk of Thrombotic Stroke and Myocardial Infarction among Users of Selected Types of Combined Oral Contraception with Ethinyl Estradiol at a Dose of 30 to 40 µg, as Compared with Nonusers, According to Duration of Use (from Lidegaard 2012).*

Duration of use	No. of person-yrs.	Thrombotic Stroke		Myocardial infarction	
		No. of events	Relative Risk (95% CI)	No. of events	Relative Risk (95% CI)
<1 year	987,564	213	1.90 (1.64–2.20)	86	1.85 (1.48–2.31)
1-4 years	992,825	194	1.55 (1.33–1.80)	108	1.99 (1.63–2.43)
>4 years	399,461	173	1.93 (1.65–2.26)	91	2.11 (1.70–2.62)

#### Atherosclerosis and Cardiovascular Events References

Bagdade JD and Subbaiah PV. Serum from Oral Contraceptive Users Stimulates Growth of Arterial Smooth Muscle Cells. *AHA Journals, Arteriosclerosis* 1982; 2(2):170–176.

Beral V. Cardiovascular disease mortality trends and oral contraceptive use in young women. *Lancet* 1976; 2:1047–1052.

Fallah S, Nouroozi V, Seifi M, Samadikuchaksaraei A, and Aghdashi EM. Influence of oral contraceptive pills on homocysteine and nitric oxide levels: as risk factors for cardiovascular disease. *J Clin Lab Anal* 2012; 26:120–123.

Hennekens CH, and MacMahon B. Oral contraceptive and myocardial infarction. *N Engl J Med* 1977; 296:1166–1167.

Hickson SS, Miles KL, McDonnell BJ, Yasmin, Cockcroft JR, Wilkinson IB, McEniery CM; ENIGMA Study Investigators. Use of the oral contraceptive pill is associated with increased large artery stiffness in young women: the ENIGMA study. *J Hypertens* 2011; 29(6):1155–1159.

Heidarzadeh Z, Asadi B, Saadatnia M, Ghorbani A, and Fatehi F. The Effect of Low-dose Combined Oral Contraceptive Pills on Brachial Artery Endothelial Function and Common Carotid Artery Intima–Media Thickness. *Journal of Stroke and Cerebrovascular Diseases* 2014; 23:675–680.

Lidegaard Ø, Løkkegaard E, Jensen A, Skovlund CW, and Keiding N. Thrombotic stroke and myocardial infarction with hormonal contraception. *N Engl J Med* 2012; 366:2257–2266.

Priest SE, Shenouda N, and MacDonald MJ. Effect of sex, menstrual cycle phase, and monophasic oral contraceptive pill use on local and central arterial stiffness in young adults. *Am J Physiol Heart Circ Physiol* 2018; 315(2):H357–H365.

Norouzi V, Seifi M, Fallah S, Korani M, and Samadikuchaksaraei A. Effect of oral contraceptive therapy on homocysteine and C-reactive protein levels in women: an observational study. *Anadolu Kardiyol Derg* 2011; 11:698–702.

Rietzschel E, De Buyzere M, De Baquer D, Langlois M, Bekaert S, Segers P, Van Damme P, Verdonck P, De Backer G, Gillebert T and ASKLEPIOS Investigators. Oral contraceptives cause major C-reactive protein rises in the female general population. *Circulation* 2018; 116:800–801.

Vessey M, Doll R, Peto R, Johnson B, and Wiggins P. A longterm follow-up study of women using different methods of contraception—an Interim report. *J Biosoc Sci* 1976; 8:373–427.

Yu A, Giannone T, Scheffler P, Doonan RJ, Egiziano G, Gomez YH, Papaioannou TG, and Daskalopoulou SS. The effect of oral contraceptive pills and the natural menstrual cycle on arterial stiffness and hemodynamicCs (CYCLIC). *J Hypertens* 2014; 32(1):100–107.

Zoler ML. Oral Contraceptive Use Linked to Arterial Plaque. *Internal Medicine News* October 15, 2007; 32.

## Conclusion

Hormonal agents have a variety of effects on various organs and organ systems which may result in a deleterious impact on women's health. The data reviewed above reflect a vast body of information which has come to light since the introduction of these agents as contraceptives over 50 years ago. While the information for patients and prescribers currently reflects many of the known side effects, others have come to light which are not adequately represented in the current prescribing information. These should be added and made obvious to patients. In one instance, that of venous thromboembolism, while the warning information is present, it is phrased in a misleading manner which misleads the patients into drawing the incorrect conclusion regarding the risks. In addition, one agent (DMPA) appears to convey a specific risk for HIV transmission which is not shared by other agents. DMPA should be considered for revoking of marketing authorization and removed from the market. The risks of depression, mood disorders, and suicide have not been adequately emphasized.

We further encourage the Agency to require the manufacturers of these agents to widely publicize these additional risks. Many millions of women are currently receiving COCs and POCs. Many millions more have been exposed to these agents at some point in their lives. They should receive updated information regarding risks which have not been conveyed, or not adequately conveyed, in the past. All women who have been exposed to COCs or POCs should be informed so that they can take this information into account as they may encounter some of these adverse effects in some cases many years after cessation of use.

## Environmental Impact

Based on data from the Guttmacher Institute, a conservative estimate of 11 million women aged 15-44 in the US take some form of hormonal contraceptive each day<sup>155</sup>. A 2015 study reports that about 21 percent of women of reproductive years are using some form of hormonal contraceptive, which equates to about 13 million women (Daniels 2015). This has resulted in a significant increase in the release of synthetic progestagens (such as levonorgestrel) and synthetic estrogens (such as ethinylestradiol [EE2]) into the aquatic environment via wastewater treatment plant discharges (Besse 2009, King 2016). EE2 is metabolized in the liver undergoing first pass metabolism, but ~6% of the administered dose appears as untransformed EE2 in the urine and ~9% in the feces (Stanczyk 2013). As noted by King (King 2016), even at low concentrations, these compounds can act as potent endocrine disruptors, affecting the growth, development, and reproduction of exposed aquatic organisms (Tyler 1998, Larsson 1999). EE2 is one of the most studied synthetic hormones in aquatic environments, for which assessments of environmental concentrations and the quantification of endocrine-related effects have been documented in a range of aquatic species (Purdom 1994, Jobling 1998, Kirby 2004, Jobling 2006). In fact, the numerous studies on the effects of EE2 on aquatic organisms have led to the derivation of a reliable predicted no-effect concentration of 0.1 ng/L for EE2 (Caldwell 2012).

In 1993, the first publication appeared which brought attention to the issue of synthetic chemicals mimicking natural estrogen in the environment (Sharpe 1993). The study pointed to environmental pollutants, which were having a deleterious effect on male fetuses in utero – endocrine disruptors like polychlorinated biphenyls, detergents, dioxins, and hormonal contraceptives. In 1995, another paper (Sumpter 1995) noted that male fish in 28 rivers across Britain were being “feminized” by pollutants. In 2002, a paper was published that focused specifically on the effects of endocrine-disrupting chemicals in the environment (Jobling 2002). They demonstrated reduced fertility in fish populations in areas downstream of effluent from sewage plants

---

<sup>155</sup> <https://www.guttmacher.org/fact-sheet/contraceptive-use-united-states>.

located along tributaries of the Thames River. In 2007, the results of a seven-year Canadian lake study were published which examined the effects of EE2 (Kidd 2007). The researchers released a quantity of EE2 equivalent to what would come into the waterways via sewage from a city of 200,000 people. They witnessed an immediate feminization and transgenerating of male fish, which resulted in the “near extinction” of the fathead minnow population (Kidd 2007). Although the minnow populations neared extinction, they rebounded as soon as the researchers stopped adding EE2 to the lake. A 2006 study from the United States Geological Survey on smallmouth bass in the Shenandoah and Monocacy Rivers found that more than 80-percent of all the male bass living in these waterways were growing eggs in their testes<sup>156</sup>.

A study was carried out of fish populations relative to the sewage treatment plants of three major Colorado cities: Denver, Boulder, and Colorado Springs (Woodling 2006). At each municipality, they set up a location just upstream from where the effluent was released, and another just downstream. The fish in the upstream locations enjoyed a balanced 1:1 female-to-male sex ratio. Downstream there were five female fish for every one male, and twenty percent of the reduced male population demonstrated intersex characteristics, such as eggs in their testes and the presence of vitellogenin, an egg yolk protein normally found only in fertile females. The consequences also appeared to ascend up the food chain in a measurable way, specifically with the feminization of trout, mink frogs and green frogs (Parke 2009). Both the predicted and the measured concentrations of EE2 in the US, including effluent of waste water treatment plants, surface water, or ground water, exceeds the predicted no-effect concentrations on fish populations (Kostich 2013).

Environmental factors have been implicated in declining fertility rates (Skakkebaek 2016). A 2017 study out of Hebrew University and Mount Sinai Medical School found that sperm counts in human men have dropped by more than 50 percent since 1973 (Levine 2017). While it has been noted that environmental exposure to individual steroidal estrogens, as well as their mixtures, are unlikely to dramatically affect endocrine signaling in humans, it is not clear whether more subtle effects are possible (Kostich 2013). More recently, environmental effects of levonorgestrel have been postulated (King 2016) but there is less hard data.

There is a clear effect of environmental EE2 on fish populations as well as species higher in the food chain such as frogs. An effect on humans is also possible.

### Environmental Impact References

Besse J-P and Garric J. Progestagens for human use, exposure and hazard assessment for the aquatic environment. *Environ Pollut* 2009; 157: 3485–3494.

Caldwell DJ, Mastrocco F, Anderson PD, L€ange R, and Sumpter JP. Predicted-no-effect concentrations for the steroid estrogens estrone, 17b-estradiol, estriol, and 17a-ethinylestradiol. *Environ Toxicol Chem* 2012; 31:1396–1406.

Daniels K, Daugherty J, Jones J, and Mosher W. Current Contraceptive Use and Variation by Selected Characteristics Among Women Aged 15–44: United States, 2011–2013. *Natl Health Stat Report* 2015; 86:1–14.

Jobling S, Nolan M, Tyler CR, Brighty G, and Sumpter JP. Widespread sexual disruption in wild fish. *Environ Sci Technol* 1998; 32:2498–2506.

---

<sup>156</sup> [https://dep.wv.gov/WWE/watershed/wqmonitoring/Documents/Potomac-Intersex/USGS\\_FishHealthReproductivelssuesPotomac\\_2006.pdf](https://dep.wv.gov/WWE/watershed/wqmonitoring/Documents/Potomac-Intersex/USGS_FishHealthReproductivelssuesPotomac_2006.pdf).

Jobling S, Williams R, Johnson A, Taylor A, Gross-Sorokin M, Nolan M, Tyler CR, van Aerle R, Santos E, and Brighty G. Predicted exposures to steroid estrogens in U.K. rivers correlate with widespread sexual disruption in wild fish populations. *Environ Health Perspect* 2006; 114:32–39.

Jobling, S, Coey S, Whitmore JG, Kime DE, Van Look KJ, McAllister BG, Beresford N, Henshaw AC, Brighty G, Tyler CR, and Sumpter JP. Wild intersex roach (*Rutilus rutilus*) have reduced fertility. *Biology of Reproduction* 2002; 67(2):515–524.

Kidd KA, Blanchfield PJ, Mills KH, Palace VP, Evans RE, Lazorchak JM, and Flick RW. Collapse of a fish population after exposure to a synthetic estrogen. *PNAS* 2007; 104(21):8897–8901.

King OC, van de Merwe JP, McDonald JA, and Leusch FD. Concentrations of levonorgestrel and ethinylestradiol in wastewater effluents: Is the progestin also cause for concern? *Environ Toxicol Chem* 2016; 35:1378–1385.

Kirby MF, Allen YT, Dyer RA, Feist SW, Katsiadaki I, Matthiessen P, Scott AP, Smith A, Stentiford GD, Thain JE, Thomas KV, Tolhurst L, and Waldock MJ. Surveys of plasma vitellogenin and intersex in male flounder (*Platichthys flesus*) as measures of endocrine disruption by estrogenic contamination in United Kingdom estuaries: Temporal trends, 1996 to 2001. *Environ Toxicol Chem* 2004; 23:748–758.

Kostich M, Flick R, and Martinson J. Comparing predicted estrogen concentrations with measurements in US waters. *Environ Pollut* 2013; 178:271–277.

Larsson DGJ, Adolfsson-Erici M, Parkkonen J, Pettersson M, Berg AH, Olsson PE, and Forlin L. Ethinylestradiol—An undesired fish contraceptive? *Aquat Toxicol* 1999; 45:91–97.

Park BJ and Kidd KA. Effects of the synthetic estrogen ethinylestradiol on early life stages of mink frogs and green frogs in the wild and in situ. *Environmental Toxicology* 2005; 24(8):2027–2036.

Purdom CE, Hardiman PA, Bye VVJ, Eno NC, Tyler CR, and Sumpter JP. Estrogenic effects of effluents from sewage treatment works. *Chem Ecol* 1994; 8:275–285.

Sharpe RM and Skakkebaek NE. Are oestrogens involved in falling sperm counts and disorders of the male reproductive tract? *Lancet* 1993; 341:1392–1395.

Skakkebaek NE, De Meyts ER, Louis GM, Toppari J, Andersson AM, Eisenberg ML, Jensen TK, Jorgensen N, Swan SH, Sapra KJ, Ziebe S, Priskorn L, and Juul A. Male reproductive disorders and fertility trends: Influences of environment and genetic susceptibility. *Physiological Reviews* 2016; 96(1):55–97.

Stanczyk FZ, Archer DF, and Bhavnani BR. Ethinyl estradiol and 17 $\beta$ -estradiol in combined oral contraceptives: pharmacokinetics, pharmacodynamics and risk assessment. *Contraception* 2013; 87:706–727.

Sumpter JP. Feminized responses in fish to environmental estrogens. *Toxicology Letters* 1995; 82-83C:737–742. 10.1016/0378-4274(95)03517-6.

Tyler CR, Jobling S, and Sumpter JP. Endocrine disruption in wildlife: A critical review of the evidence. *Crit Rev Toxicol* 1998; 28:319–319.



Woodling, J, Lopez E, Maldonado T, Norris D, and Vajda A. Intersex and other reproductive disruption of fish in wastewater effluent dominated Colorado streams. *Comparative Biochemistry and Physiology C Toxicology and Pharmacology* 2006; 144 (1):10–15.

## Economic Impact

For the diseases noted below, in some cases we have calculated the estimated economic impact taking into account those who are currently using COCs and those who have ever used COCs. According to the CDC<sup>157</sup> 15.9% of women aged 15–44 in the US use “the pill.” There are 61 million US women of reproductive age (15–44)<sup>158</sup>. This yields 9,699,000 women in the USA currently on COCs. Note that this is a low estimate as it does not include women using intravaginal and transdermal formulations and is lower than the estimate by Daniels (Daniels 2015).

According to the National Survey of Family Growth<sup>159</sup>, 79.3% of women surveyed from 2011–2015 have ever used “the pill.” This is down from 81.9% in the 2006–2010 survey and 82.3% in the 2002 survey. The lower number for “ever use” of 79.3% is used in subsequent calculations. According to the 2010 census (Howden 2011), there were 156,964,212 women in the US, of whom 24% were under 18 years of age. Thus, there were 119,292,801 women 18 years of age or older. This implies that  $119,292,801 \times 0.793 = 94,599,191$  women in the USA have ever used the pill. As noted above, this does not include women using intravaginal and transdermal formulations.

The numbers 9,699,000 for current use and 94,599,191 for ever use of COCs were used in some of these calculations. In other cases, the census data for specific age groups was used if they were the groups most likely to be impacted by current or recent use of COCs.

For progesterone-only contraceptives (POCs), the National Survey of Family Growth<sup>160</sup>, notes that 25.4% of women aged 15–44 in 2011–2015 have ever used “3-month injectable (Depo-Provera™).” This is up from 23.2% in 2006–2010 and 16.8% in 2002. For a conservative estimate, we will use the lowest of these numbers (16.8% or 20,041,191 women) who have ever used POCs. This would not include POCs administered by other routes and is thus a conservative estimate.

## HIV Costs

According to the CDC<sup>161</sup>, an estimated 255,900 women were living with HIV at the end of 2014. Of these it is estimated 87% were via sexual contact (this proportion was relatively stable from 2011–2016; CDC HIV Surveillance Table 1a). Annual medical cost estimates for HIV-infected persons, adjusted for age, sex, race/ethnicity, and transmission risk group, were from the HIV Research Network (range \$1,854–\$4,545/month) and for HIV-uninfected persons were from the Medical Expenditure Panel Survey (range \$73–\$628/month) (Schackman 2015). Using this information along with the prevalence of DMPA use of 16.8%, this suggests an annual cost of treatment for HIV infection due to DMPA use of ~\$157–573 million (Table 16).

---

<sup>157</sup> <https://www.cdc.gov/nchs/fastats/contraceptive.htm>.

<sup>158</sup> <https://www.cdc.gov/nchs/data/nhsr/nhsr086.pdf>.

<sup>159</sup> [https://www.cdc.gov/nchs/nsfg/key\\_statistics/c.htm#everused](https://www.cdc.gov/nchs/nsfg/key_statistics/c.htm#everused).

<sup>160</sup> [https://www.cdc.gov/nchs/nsfg/key\\_statistics/c.htm#everused](https://www.cdc.gov/nchs/nsfg/key_statistics/c.htm#everused).

<sup>161</sup> <https://www.cdc.gov/hiv/group/gender/women/index.html>.

Table 16 – Estimated Economic Impact of DMPA due to Increased Prevalence of HIV Infection

Women with HIV	255,900
Sexual transmission	87%
Cases due to sexual transmission	222,633
Ever use of DMPA	16.80%
Women with HIV with DMPA use	37,402
RR of HIV with DMPA use	1.4
Adjusted estimate →	26,716
Excess cases →	10,686
Highest estimated individual annual costs →	\$53,664
Lowest estimated individual annual costs →	\$14,712
Highest estimated total annual costs →	\$573,474,111
Lowest estimated total annual costs →	\$157,218,081

### Breast Cancer

A recent study in the US (Blumen 2016) notes, “The average costs per patient allowed by the insurance company in the year after diagnosis were \$60,637, \$82,121, \$129,387, and \$134,682 for disease stage 0, I/II, III, and IV, respectively. The average costs allowed per patient in the 24 months after the index diagnosis were \$71,909, \$97,066, \$159,442, and \$182,655 for disease stage 0, I/II, III, and IV, respectively.” For all patients, they note that the average cost for the first 12 months following diagnosis is \$85,772, and for the second 12 months is \$22,127 with a total of \$103,735 for the 24 months following diagnosis. For these calculations we will use the first-year costs to estimate costs for incident cases among current users of COCs and will use the second-year cost to approximate the average annual cost of care for a patient diagnosed with breast cancer. According to the NIH SEER statistics<sup>162</sup>, the incidence of breast cancer is 126.0 per 100,000 person-years. Approximately 12.4 percent of women will be diagnosed with female breast cancer at some point during their lifetime. According to the best epidemiology studies noted in Table 3 (Mørch 2017; Heikkinen 2016, Lund 2007), and the best meta-analysis in Table 5 (Kahlenborn 2006) the relative risk of ever use of COCs for the development of breast cancer is 1.19–1.37. Based on this information, the estimated increase in cost from use of COCs due to incident cases of breast cancer is between \$199 million and \$387 million (Table 17).

Table 17 – Estimated Economic Impact of COCs due to Increased Incidence of Breast Cancer

Women of reproductive age	Number on the pill	Incidence		
61,000,000	9,699,000	0.00126		
Estimated women on the pill at risk →		12,221		
Adjusted estimate of cases →		14,543	1.19	Low RR
Adjusted estimate of cases →		16,742	1.37	High RR
Excess cases →		2,322	Low RR	
Excess cases →		4,522	High RR	
Annual cost per patient of breast cancer →		\$85,772		
Estimated annual costs →		\$199,157,489	Low RR	
Estimated annual costs →		\$387,833,005	High RR	

<sup>162</sup> <https://seer.cancer.gov/statfacts/html/breast.html>.

To evaluate the impact of “ever use” of COCs on prevalent breast cancer, we noted that the best meta-analysis (Kahlenborn 2006) showed a 1.19 odds ratio of breast cancer with COCs. According to the SEER statistics, there are currently 3,418,124 prevalent cases of breast cancer in the USA. The estimated increase in cost from treatment of the excess cases of breast cancer is estimated to be ~\$9.6 billion annually (Table 18).

*Table 18 – Estimated Economic Impact of COCs due to Increased Prevalence of Breast Cancer*

Prevalent cases of breast cancer	Ever use of COCs	Breast cancer ever users		
3,418,124	79.3%	2,710,572		
Adjusted estimate of cases if no use of COCs →		2,277,792	1.19	RR
Excess cases →		432,780		
Annual cost per patient of breast cancer →				\$22,127
Estimated total costs →				\$9,576,133,158

### Cervical Cancer

A recent study in Canada (Pendrith 2016) on the costs of invasive cervical cancer treatment noted: “The mean overall medical care cost was \$39,187 [standard error (se): \$1,327] in the 1st year after diagnosis. ... At 5 years after diagnosis, the mean overall unadjusted cost was \$63,131 (se: \$3,131), and the cost adjusted for censoring was \$68,745 (se: \$2,963).” For these calculations we will assume a cost of \$39,187 annually for incident cases and \$13,749 (= \$68,745/5) annually for prevalent cases of invasive cervical cancer. According to the NIH SEER statistics<sup>163</sup>, the incidence of invasive cervical cancer is 7.4 per 100,000 person-years. According to the American Cancer Society<sup>164</sup>, it is estimated that 13,170 women will be diagnosed with invasive cervical cancer in the USA in 2019. In 2015, there were an estimated 257,524 women living with invasive cervical cancer in the United States. According to the best epidemiology studies noted in Table 6 (Roura 2016) the relative risk of ever use of COCs for the development of invasive cervical cancer is 1.6 and the RR for current use is 2.2. Based on this information, the estimated increase in cost from use of COCs due to incident cases of cervical cancer is ~\$33 million (Table 19).

*Table 19 – Estimated Economic Impact of COCs due to Increased Incidence of Cervical Cancer*

Women of reproductive age	Number on the pill	Incidence		
61,000,000	9,699,000	0.000074		
Estimated women on the pill at risk →		718		
Adjusted estimate of cases →		1,579	2.2	RR
Excess cases →		861		
Annual cost per patient of cervical cancer →		\$39,187		
Estimated annual costs →		\$33,750,635		

To evaluate the impact of “ever use” of COCs on prevalent cervical cancer, we noted that the best study (Roura 2016) showed a 1.6 relative risk of cervical cancer with COCs. According to the SEER statistics, there are currently 257,524 prevalent cases of cervical cancer in the USA. The estimated increase in cost from treatment of the excess cases of cervical cancer is estimated to be ~\$1 billion annually (Table 20).

<sup>163</sup> <https://seer.cancer.gov/statfacts/html/cervix.html>.

<sup>164</sup> <https://www.cancer.org/cancer/cervical-cancer/about/key-statistics.html>.

Table 20 – Estimated Economic Impact of COCs due to Increased Prevalence of Cervical Cancer

Prevalent cases of breast cancer	Ever use of COCs	Cervical cancer ever users		
3,418,124	79.3%	257,524		
Adjusted estimate of cases if no use of COCs →		204,217	1.6	RR
	Excess cases →	76,581		
Annual cost per patient of cervical cancer →			\$13,749	
	Estimated total costs →		\$1,052,914,912	

### Crohn’s Disease

A recent study in the US (Rao 2018) estimated the 5-year cost of the treatment of Crohn’s disease as \$116,838 per patient (interquartile range of \$45,643–\$240,398; annual cost \$23,368). This was higher with worsening disease activity. According to the Centers for Disease Control (CDC), the incidence of Crohn’s disease is 3.1 to 14.6 cases per 100,000 person-years<sup>165</sup>. According to the best epidemiology studies noted in Table 7 (Khalili 2013; García Rodríguez 2005), and the best meta-analysis (Cornish 2008), the relative risk of current COC use is 1.46–2.82 for the development of Crohn’s disease. Based on this information, the estimated increase in cost just from treatment of the excess cases of Crohn’s disease, only looking at current use and not past use of COCs, is between \$3 million and \$60 million annually (Table 21).

Table 21 – Estimated Economic Impact of COCs due to Increased Incidence of Crohn’s Disease

Women of reproductive age	Number on the pill	Low incidence	High incidence		
61,000,000	9,699,000	0.000031	0.000146		
Estimated women on the pill at risk →		301	1,416		
Adjusted estimate →		439	2,067	1.46	Low RR
Adjusted estimate →		848	3,993	2.82	High RR
Excess cases →		138	651	Low RR	
Excess cases →		547	2,577	High RR	
Annual cost per patient of Crohn’s disease →		\$23,368			
Estimated annual costs →		\$3,231,920	\$15,221,300	Low RR	
Estimated annual costs →		\$12,787,162	\$60,223,406	High RR	

To evaluate the impact of “ever use” of COCs, we noted that the best cohort study (Khalili 2013) and meta-analysis (Cornish 2008) showed a 1.43 and 1.44 relative risk of Crohn’s disease. According to the Centers for Disease Control (CDC), the prevalence of Crohn’s disease in adults is 201 cases per 100,000 person-years<sup>166</sup>. Taking the lower number of 1.43, the estimated increase in cost from treatment of the excess cases of Crohn’s disease due to COC use is approximately \$1.9 billion annually (Table 22).

Table 22 – Estimated Economic Impact of COCs due to Increased Prevalence of Crohn’s Disease

Women ≥ 18 in 2010 Census	Ever use of COCs	Prevalence	
---------------------------	------------------	------------	--

<sup>165</sup> <https://www.cdc.gov/ibd/IBD-epidemiology.htm>.

<sup>166</sup> <https://www.cdc.gov/ibd/IBD-epidemiology.htm>.

119,292,801	94,599,191	0.000201		
Estimated women on the pill at risk →		190,144		
Adjusted estimate →		271,906	1.44	RR
Excess cases →		81,762	1.44	RR
Annual cost per patient of Crohn's disease →		\$23,368		
Estimated total costs →		\$1,910,583,605	1.44	RR

### Ulcerative Colitis

A recent study in the US (Cohen 2015) noted that compared with controls, patients with UC had higher adjusted total direct (\$15,548 vs \$4812) and indirect costs (\$4125 vs \$1961) annually. This implies a total annual increase in cost of ~\$12,900 for UC. This was higher with worsening disease activity. According to the Centers for Disease Control (CDC), the incidence of UC is 2.2 to 14.3 cases per 100,000 person-years<sup>167</sup>. According to the best epidemiology studies noted in Table 8 (Khalili 2013; García Rodríguez 2005), and the best meta-analysis (Cornish 2008) the relative risk of current COC use 1.22–1.58 for the development of UC. Based on this information, the estimated increase in cost just from treatment of the excess cases of UC, only looking at current use and not past use of COCs is between \$605,000 and \$10 million per year (Table 23).

*Table 23 – Estimated Economic Impact of COCs due to Increased Incidence of Ulcerative Colitis*

Women of reproductive age	Number on the pill	Low incidence	High incidence		
61,000,000	9,699,000	0.000022	0.000143		
Estimated women on the pill at risk →		213	1,387		
Adjusted estimate →		260	1,692	1.22	Low RR
Adjusted estimate →		337	2,191	1.58	High RR
Excess cases →		47	305	Low RR	
Excess cases →		124	804	High RR	
Annual cost per patient of ulcerative colitis →		\$12,900			
Estimated annual costs →		\$605,567	\$3,936,184	Low RR	
Estimated annual costs →		\$1,596,494	\$10,377,212	High RR	

To evaluate the impact of “ever use” of COCs, we noted that the best cohort study (Khalili 2013) showed a 1.18 relative risk of UC. The estimated increase in cost of the excess cases of UC due to use of COCs is approximately \$522 million annually (Table 24).

*Table 24 – Estimated Economic Impact of COCs due to Increased Prevalence of Ulcerative Colitis*

Women ≥ 18 in 2010 Census	Ever use of COCs	Prevalence		
119,292,801	94,599,191	0.000238		
Estimated women on the pill at risk →		225,146		
Adjusted estimate →		265,672	1.18	RR
Excess cases →		40,526	1.18	RR
Annual cost per patient of ulcerative colitis →		\$12,900		
Estimated total costs →		\$522,789,187	1.18	RR

<sup>167</sup> <https://www.cdc.gov/ibd/IBD-epidemiology.htm>.

## Systemic Lupus Erythematosus

A recent study in the US (Chen 2015) noted that mean total health care costs were \$21,535 among all SLE patients over the 1-year study period. According to the Centers for Disease Control (CDC), the incidence of SLE is 6.5–10.6 cases per 100,000 women-years<sup>168</sup>. In terms of prevalence, “A conservative estimate suggests a prevalence of 161,000 with definite SLE and 322,000 with definite or probable SLE.” According to the best epidemiology studies noted in Table 9 that evaluated current use of COCs (Bernier 2009), the relative risk of current COC use is 1.45 – 2.52 for the development of SLE. Based on this information, the estimated increase in cost just from treatment of the excess cases of SLE, only looking at current use and not past use of COCs, is \$6.1 million to \$33.6 million annually (Table 25).

*Table 25 – Estimated Economic Impact of COCs due to Increased Incidence of Systemic Lupus Erythematosus.*

Women of reproductive age	Number on the pill	Low incidence	High incidence		
61,000,000	9,699,000	0.000065	0.0001065		
Estimated women on the pill at risk →		630	1,028		
Adjusted estimate →		914	1,491	1.45	Low RR
Adjusted estimate →		1,589	2,591	2.52	High RR
Excess cases →		284	463	Low RR	
Excess cases →		958	1,563	High RR	
Annual cost per patient of SLE →		\$21,535			
Estimated annual costs →		\$6,109,388	\$9,963,002	Low RR	
Estimated annual costs →		\$20,636,155	\$33,652,807	High RR	

To evaluate the impact of “ever use” of COCs, we noted that the best cohort studies (Costenbader 2007; Bernier 2009) showed a relative risk of SLE 1.19–2.3. The estimated increase in cost of the excess cases of SLE due to use of COCs is approximately \$439 million–\$1.55 billion annually (Table 26).

*Table 26 – Estimated Economic Impact of COCs due to Increased Prevalence of Systemic Lupus Erythematosus.*

Women ≥ 18 in 2010 Census	Ever use of COCs	Prevalence			
119,292,801	94,599,191	161,000			
Estimated women on the pill at risk →		127,673			
Adjusted estimate →		107,288	1.19	Low RR	
Adjusted estimate →		55,510	2.3	High RR	
Excess cases →		20,385	1.19	Low RR	
Excess cases →		72,163	2.3	High RR	
Annual cost per patient of SLE →		\$21,535			
Estimated total costs →		\$438,985,908	1.19	Low RR	
Estimated total costs →		\$1,554,030,205	2.3	High RR	

## Depression

The most reliable study (Skovlund 2016) indicated a 1.1 RR for depression with COCs and a 1.2 RR with POCs. This study evaluated women aged 15-34 and then followed them for a mean of 5 years. According to the information from Brody (Brody 2018), the prevalence of depression in women aged 20-39 is 10.1%. An analysis

<sup>168</sup> <https://www.cdc.gov/lupus/facts/detailed.html>.

of medical claims conducted by insurer Blue Cross Blue Shield (Blue Cross Blue Shield 2018) found that “in 2016, Blue Cross plans spent \$10,673 on those diagnosed with ‘major depression’ compared to \$4,283 on those without a depression diagnosis.” With this information, and noting from the census data (Howden 2011) that there are ~52 million women aged 15-39, we calculate that the excess annual cost of depression attributable to COCs is ~\$2.4 billion (Table 27) and from POCs is ~\$937 million (Table 28).

*Table 27 – Estimated Economic Impact of COCs due to Increased Prevalence of Depression*

Women aged 15-39	51,877,977
Percent with depression	10.1%
Women aged 15-39 with depression	5,239,675.68
Ever use of COCs	79.30%
15-39 y.o. COC users with depression	4,155,063
RR of depression with COC use	1.1
Adjusted estimate →	3,777,330
Excess cases →	377,733
Estimated individual annual costs →	\$6,390
Estimated total annual costs →	\$2,413,713,761

*Table 28 – Estimated Economic Impact of POCs due to Increased Prevalence of Depression*

Women aged 15-39	51,877,977
Percent with depression	10.1%
Women aged 15-39 with depression	5,239,675.68
Ever use of POCs	16.80%
15-39 y.o. COC users with depression	880,266
RR of depression with POC use	1.2
Adjusted estimate →	733,555
Excess cases →	146,711
Estimated individual annual costs →	\$6,390
Estimated total annual costs →	\$937,482,772

## Multiple Sclerosis

As the most rigorous cohort studies did not show an increase in the risk of developing multiple sclerosis a rigorous cost analysis was not performed. However, using the information from the best case-control study (Hellwig 2016), an increased odds ratio of 1.51 was noted. If this is assumed to be accurate, this can be used along with a study of total MS costs from 1997-2013 (Chen 2017). They noted that, “The total charges on managing MS range from \$161 million in 1997 to \$755 million in 2013.” Conservatively assuming steady costs since 2013, we can calculate that 79.3% of those costs were incurred by women who were “ever users” of



COCs. This yields \$598,715,000. If these women had not used COCs there would have been a proportionate reduction in costs of \$202,215,000 ( $\$598,715,000 - (\$598,715,000/1.51)$ ).

### Interstitial Cystitis

According to one recent paper (Tung 2017) on average, having interstitial cystitis was associated with \$7,223 higher total health care costs annually than not having IC. The prevalence of interstitial cystitis has been estimated at 2.7% using a high specificity definition (McLennan 2014) while another study in a managed care population (Clemens 2005) indicated (depending on the definition) a prevalence between 45 and 197 per 100,000 women. Using the most conservative estimate (Champaneria 2015) “ever use” of COCs is associated with an OR of 2.31 for interstitial cystitis. Assuming 61 million women of reproductive age, with a 79.3% of exposure to COCs, this suggests ~11,500 excess cases (using a prevalence of interstitial cystitis of 45/100,000) to ~50,500 (using a prevalence of interstitial cystitis of 197/100,000). This yields an annual cost of \$83–\$365 million (Table 29).

*Table 29 – Estimated Annual Economic Impact of COCs due to Increased Prevalence of Interstitial Cystitis*

Low prevalence of interstitial cystitis	0.00045
High prevalence of interstitial cystitis	0.00197
Women of reproductive age	61,000,000
Number with ever use of the pill	48,373,000
# of Women with interstitial cystitis low prevalence	21,768
# of Women with interstitial cystitis high prevalence	95,295
OR	2.13
Excess cases of interstitial cystitis low prevalence	11,548
Excess cases of interstitial cystitis high prevalence	50,555
Annual cost	\$7,223
Annual cost of interstitial cystitis low prevalence	\$83,412,664
Annual cost of interstitial cystitis high prevalence	\$365,162,106

### Osteoporotic Bone Fracture Risk

According to a recent review (Ballane 2017), in North America the incidence of osteoporotic vertebral fractures is 837 to 1,083 cases per 100,000 women per year (mean of 960 per 100,000 per year) as standardized to 2015. The annual excess cost of care for women with osteoporotic vertebral fractures was estimated to be \$11,655 per year (Kilgore 2009). Using the most relevant relative risk of 1.07 (Barad 2005), this implies an annual cost of ~\$308 million dollars in the US from COC use (Table 30).

*Table 30 – Estimated Economic Impact of COCs due to Increased Annual Incidence of Vertebral Fractures*

Women ≥ 50 in 2010 Census	Ever use of COCs	Incidence of osteoporotic vertebral fractures		
53,151,456	42,149,105	0.0096		
Estimated women on the pill with Fx →		404,631		
Adjusted estimate →		378,160	1.07	RR
Excess cases →		26,471		
Annual cost per patient of osteoporotic vertebral fractures →		\$11,655		



Estimated total annual costs →	\$308,521,992
--------------------------------	---------------

The best cohort study on fracture risk with progesterone-only contraceptives (POCs) showed a RR of 1.51 for ever use of DMPA (Lanza 2013), the most widely used POC. Assuming 16.8% of women have used POCs this yields an annual cost of ~\$290 million dollars in the US from POC use (Table 31).

Table 31 – Estimated Economic Impact of POCs due to Increased Annual Incidence of Vertebral Fractures

Women ≥ 50 in 2010 Census	Ever use of POCs	Incidence of osteoporotic vertebral fractures		
53,151,456	8,929,445	0.0096		
Estimated women on the pill with Fx →		85,723		
Adjusted estimate →		60,796	1.41	RR
Excess cases →		24,926		
Annual cost per patient of osteoporotic vertebral fractures →		\$11,655		
Estimated total annual costs →		\$290,517,770		

### Body Mass

The costs of the effects on body mass were not calculated, but these effects are contributory to atherosclerosis and cardiovascular events, which are discussed below.

### Urogenital Effects

The medical and societal costs of the urogenital effects of hormonal contraceptives were not calculated as, although there are measurable costs, they are not felt to be significant.

### Venous Thromboembolism, Atherosclerosis and Cardiovascular Disease

About 1 in every 4 female deaths is due to heart disease; it is the leading cause of death for women in the U.S.<sup>169</sup> A review of recent population studies revealed that the overall prevalence of Peripheral Arterial Disease (PAD) for women is 15.6% (compared to 13.4% for men).<sup>170</sup> In 2008, coronary heart disease was prevalent in 7.5 million women.<sup>171</sup> The total mean direct medical costs for cardiovascular disease (CVD) is \$18,953 annually (Nichols 2010). Using the median relative risk of the most popular birth control brands, the RR is 1.8 (Table 32).

<sup>169</sup> [https://www.cdc.gov/dhds/data\\_statistics/fact\\_sheets/fs\\_women\\_heart.htm](https://www.cdc.gov/dhds/data_statistics/fact_sheets/fs_women_heart.htm).

<sup>170</sup> [https://www.medscape.org/viewarticle/711179\\_2](https://www.medscape.org/viewarticle/711179_2).

<sup>171</sup> <https://www.healthline.com/health/heart-disease/women-statistics-facts#1>.

Table 32 – Estimated Economic Impact of COCs due to Increased Incidence of Cardiovascular Disease

Coronary heart disease	Ever use of COCs	CHD in ever users		
7,500,000	79.3%	5,947,500		
Adjusted estimate of cases if no use of COCs →		3,304,167	1.8	RR
Excess cases →		2,643,333		
Annual cost per patient for CVD →		\$18,953		
Estimated total costs →		\$50,099,090,349		

A more conservative estimate would assume that the increased risk for cardiovascular disease is limited to women aged 15–49 years, which was the group studied by Lidegaard (Lidegaard 2012). According to the US Census in 2010, population is broken down by age group (Howden 2011). The rate of cardiovascular events is similarly broken down by Lidegaard (Lidegaard 2012). Thus, the number of cases by age group is shown in Table 33.

Table 33 – Cardiovascular Events in Women by Age Group

Census data		Events per 100,000 person-years (Lidegaard 2012)		Events per year	
Age group	Number of women	Myocardial infarction	Stroke	Myocardial infarction #	Stroke #
15 to 19 years	10,736,677	0.4	3.4	43	365
20 to 24 years	10,571,823	0.7	5.6	74	592
25 to 29 years	10,466,258	2.2	10.5	230	1,099
30 to 34 years	9,965,599	5	15.4	498	1,535
35 to 39 years	10,137,620	12.2	23.3	1,237	2,362
40 to 44 years	10,496,987	25.4	39.2	2,666	4,115
45 to 49 years	11,499,506	38.2	64.4	4,393	7,406
Total number of events per year				9,141	17,473

Using these estimates, with the annual cost of care for cardiovascular disease and the relative risk noted above, this calculates to ~\$61 million in excess costs for myocardial infarctions and ~\$117 million in excess costs for strokes (Table 34).

Table 34 – Cost of Cardiovascular Events in Women Attributable to COC use.

	Myocardial infarction	Stroke		
Total events per year	9,141	17,473		
Ever use of the pill	79.30%			
# with events on COCs	7,249	13,856	1.8	RR Ever Use
Adjusted estimate →	4,027	7,697.96		
Excess cases →	3,222	6,158		
Estimated annual costs →	\$18,953	\$18,953		
Estimated Excess annual costs →	\$61,062,935	\$116,719,504		

#### Economic Costs References

Ballane G, Cauley JA, Luckey MM, and El-Hajj Fuleihan G. Worldwide prevalence and incidence of osteoporotic vertebral fractures. *Osteoporos Int* 2017; 28(5):1531–1542.

Barad D, Kooperberg C, Wactawski-Wende J, Liu J, Hendrix SL, and Watts NB. Prior oral contraception and postmenopausal fracture: a Womens' Health Initiative observational cohort study. *Fertility and Sterility* 2005; 84:374–383.

Bernier MO, Mikaeloff Y, Hudson M, and Suissa S. Combined oral contraceptive use and the risk of systemic lupus erythematosus. *Arthritis and Rheumatism* 2009; 61:476–481.

Blue Cross Blue Shield. Major Depression: The Impact on Overall Health. Blue Cross Blue Shield, The Health of America Report®, 2018.

Blumen H, Fitch K, and Polkus V. Comparison of Treatment Costs for Breast Cancer, by Tumor Stage and Type of Service. *Am Health Drug Benefits* 2016; 9(1):23–32.

Brody DJ, Pratt LA, and Hughes JP. Prevalence of Depression Among Adults Aged 20 and Over: United States, 2013-2016. *NCHS Data Brief* 2018; 303:1–8.

Champaneria R, D'Andrea RM, and Latthe PM. Hormonal contraception and pelvic floor dysfunction: a systematic review. *Int Urogynecol J* 2016; 27:709–722.

Chen AY, Chonghasawat AO, and Leadholm KL. Multiple sclerosis: Frequency, cost, and economic burden in the United States. *J Clin Neurosci* 2017; 45:180–186.

Chen SY, Choi CB, Li Q, Yeh WS, Lee YC, Kao AH, and Liang MH. Glucocorticoid Use in Patients With Systemic Lupus Erythematosus: Association Between Dose and Health Care Utilization and Costs. *Arthritis Care Res (Hoboken)*. 2015; 67(8):1086–1094.

Clemens JQ, Meenan RT, Rosetti MC, Gao SY, and Calhoun EA. Prevalence and incidence of interstitial cystitis in a managed care population. *J Urol*. 2005; 173(1):98–102; discussion 102.

Cohen R, Skup M, Ozbay AB, Rizzo J, Yang M, Diener M, and Chao J. Direct and indirect healthcare resource utilization and costs associated with ulcerative colitis in a privately-insured employed population in the US. *J Med Econ* 2015; 18(6):447–456.

Cornish JA, Tan E, Simillis C, Clark SK, Teare J, and Tekkis PP. The risk of oral contraceptives in the etiology of inflammatory bowel disease: a meta-analysis. *American Journal of Gastroenterology* 2008; 103:2394–2400.

Costenbader KH, Feskanich D, Stampfer MJ, and Karlson EW. Reproductive and menopausal factors and risk of systemic lupus erythematosus in women. *Arthritis and Rheumatism* 2007; 56:1251-1262.

Daniels K, Daugherty J, Jones J, and Mosher W. Current Contraceptive Use and Variation by Selected Characteristics Among Women Aged 15-44: United States, 2011-2013. *Natl Health Stat Report* 2015; (86):1–14.

García Rodríguez LA, González-Pérez A, Johansson S, and Wallander MA. Risk factors for inflammatory bowel disease in the general population. *Alimentary Pharmacology & Therapeutics* 2005; 22:309–315.

Heikkinen S, Koskenvuo M, Malila N, Sarkeala T, Pukkala E, and Pitkaniemi J. Use of exogenous hormones and the risk of breast cancer: results from self-reported survey data with validity assessment. *Cancer Causes Control* 2016; 27:249–258.

Hellwig K, Chen LH, Stanczyk FZ, and Langer-Gould AM. Oral Contraceptives and Multiple Sclerosis/Clinically Isolated Syndrome Susceptibility. *PLoS One* 2016; 11:e0149094. Doi: 10.1371/journal.pone.0149094.

Howden LM, and Meyer JA. Age and Sex Composition. *2010 Census Briefs*. U.S. Department of Commerce Economics and Statistics Administration, U.S. Census Bureau. 2011.

Kahlenborn C, Modugno F, Potter DM, and Severs WB. Oral Contraceptive Use as a Risk Factor for Premenopausal Breast Cancer: A Meta-analysis. *Mayo Clin Proc* 2006; 81(10):1290–1302.

Khalili H, Higuchi LM, Ananthakrishnan AN, Richter JM, Feskanich D, Fuchs CS, and Chan AT. Oral contraceptives, reproductive factors and risk of inflammatory bowel disease. *Gut* 2013; 62:1153–1159.

Kilgore ML, Morrissey MA, Becker DJ, Gary LC, Curtis JR, Saag KG, Yun H, Matthews R, Smith W, Taylor A, Arora T, and Delzell E. Health care expenditures associated with skeletal fractures among Medicare beneficiaries, 1999-2005. *J Bone Miner Res* 2009; 24(12):2050–2055.

Lanza LL, McQuay LJ, Rothman KJ, Bone HG, Kaunitz AM, Harel Z, Ataher Q, Ross D, Arena PL, and Wolter KD. Use of depot medroxyprogesterone acetate contraception and incidence of bone fracture. *Obstetrics and Gynecology* 2013; 121(3):593–600.

Lidegaard Ø, Nielsen LH, Skovlund CW, and Løkkegaard E. Venous thrombosis in users of non-oral hormonal contraception: follow-up study, Denmark 2001-10. *British Medical Journal* 2012; 344:e2990.

Lund E, Bakken K, Dumeaux V, Andersen V, and Kumle M. Hormone replacement therapy and breast cancer in former users of oral contraceptives—The Norwegian Women and Cancer study. *Int J Cancer* 2007; 121:645-648.

McLennan MT. Interstitial cystitis: epidemiology, pathophysiology, and clinical presentation. *Obstet Gynecol Clin North Am* 2014; 41(3):385-395.

Mørch LS, Skovlund CW, Hannaford PC, Iversen L, Fielding S, and Lidegaard Ø. Contemporary hormonal contraception and the risk of breast cancer. *N Engl J Med* 2017; 377:2228–2239.

Nichols GA, Bell TJ, Pedula KL, and Rosetti MO. Medical care costs among patients with established cardiovascular disease. *The American Journal of Managed Care* 2010; 16(3):e86–e93.

Pendrith C, Thind A, Zaric GS, and Sarma S. Costs of cervical cancer treatment: population-based estimates from Ontario. *Curr Oncol* 2016; 23(2):e109–15.

Rao BB, Click BH, Koutroubakis IE, Ramos Rivers C, Regueiro M, Swoger J, Schwartz M, Hashash J, Barrie A, Dunn MA, and Binion DG. The Cost of Crohn's disease: varied healthcare expenditure patterns across distinct disease trajectories. *Inflamm Bowel Dis* 2017; 23(1):107–115.

Roura E, Travier N, Waterboer T, de Sanjosé S, Bosch FX, Pawlita M, Pala V, Weiderpass E, Margall N, Dillner J, Gram IT, Tjønneland A, Munk C, Palli D, Khaw KT, Overvad K, Clavel-Chapelon F, Mesrine S, Fournier A, Fortner RT, Ose J, Steffen A, Trichopoulou A, Lagiou P, Orfanos P, Masala G, Tumino R, Sacerdote C, Polidoro S, Mattiello A, Lund E, Peeters PH, Bueno-de-Mesquita HB, Quirós JR, Sánchez MJ, Navarro C, Barricarte A, Larrañaga N, Ekström J, Lindquist D, Idahl A, Travis RC, Merritt MA, Gunter MJ, Rinaldi S, Tommasino M, Franceschi S, Riboli E, and Castellsagué X. The Influence of Hormonal Factors on the Risk of Developing Cervical Cancer and Pre-Cancer: Results from the EPIC Cohort. *PLoS One* 2016; 11:e0147029.

Schackman BR, Fleishman JA, Su AE, Berkowitz BK, Moore RD, Walensky RP, Becker JE, Voss C, Paltiel AD, Weinstein MC, Freedberg KA, Gebo KA, and Losina E. The lifetime medical cost savings from preventing HIV in the United States. *Med Care* 2015; 53(4):293–301.

Skovlund CW, Mørch LS, Kessling LV, and Lidegaard O. Association of Hormonal Contraception with Depression. *JAMA Psychiatry* 2016; 73(11):1154–1162.

Tung A, Hepp Z, Bansal A, and Devine EB. Characterizing Health Care Utilization, Direct Costs, and Comorbidities Associated with Interstitial Cystitis: A Retrospective Claims Analysis. *J Manag Care Spec Pharm* 2017; 23(4):474–482.



## Certification

We certify that this petition contains all relevant information, including any that may be unfavorable to the petition, that we were able to obtain.

### Main Representative:

*William V. Williams*

Date: *2019 May 09*

William V. Williams, M.D.  
Contraceptive Study Group  
620 South Eagle Road  
Havertown, PA 19083  
Phone: (302) 290-9017  
E-Mail: william.v.williams@verizon.net

### Other Members of the Group:

Joel Brind, Ph.D.

Michael D. Manhart, Ph.D.

Hanna Klaus, M.D.

Angela Lanfranchi, M.D.

Gerard Migeon

Michael Gaskins

Elvis Seman, M.D.

Lester Ruppertsberger, D.O.

Kathleen Raviele, M.D.

Dear Food and Drug Administration,

The following organizations and individuals strongly encourage the FDA to carefully review the Citizens Petition on Hormonal Contraceptives.

<u>Name</u>	<u>Email</u>	<u>City and State of Residence</u>
<b>Organizations</b>		
Catholic Medical Association	<a href="mailto:info@cathmed.org">info@cathmed.org</a>	Bala Cynwyd, PA
Care Net	<a href="mailto:vdicaro@care-net.org">vdicaro@care-net.org</a>	Lansdowne, VA
Coalition on Abortion Breast Cancer, signed by Jo Ann Gerling	<a href="mailto:rjgerling@aol.com">rjgerling@aol.com</a>	New Brunswick, NJ
Couple to Couple League, signed by Executive Director Chris Reynolds	<a href="mailto:creynolds@ccli.org">creynolds@ccli.org</a>	Cincinnati, OH
Family of the Americas, signed by Mercedes Wilson	<a href="mailto:fafmercedes@yahoo.com">fafmercedes@yahoo.com</a>	Lothian, MD
Fertility Appreciation Collaborative to Teach the Science (FACTS), signed by Executive Director Marguerite Duane	<a href="mailto:drduane@factsaboutfertility.org">drduane@factsaboutfertility.org</a>	Washington, DC
Good Counsel, Inc., signed by President and CEO Christopher Bell	<a href="mailto:cxbell@aol.com">cxbell@aol.com</a>	Hoboken, NJ
Human Life Alliance, signed by Executive Director Joe Langfeld	<a href="mailto:jlangfeld@humanlife.org">jlangfeld@humanlife.org</a>	Minneapolis, MN
Institute for Natural Family Planning, College of Nursing, Marquette University, signed by Dr. Richard Fehring	<a href="mailto:richard.fehring@marquette.edu">richard.fehring@marquette.edu</a>	Milwaukee, WI
National Catholic Bioethics Center, signed by President John Haas	<a href="mailto:jhaas@ncbcenter.org">jhaas@ncbcenter.org</a>	Philadelphia, PA
Northwest Family Services	<a href="mailto:kdetloff@nwfs.org">kdetloff@nwfs.org</a>	Portland, OR
Population Research Institute, signed by	<a href="mailto:steve@pop.org">steve@pop.org</a>	Front Royal, PA

President Steven Mosher		
St. Augustine Foundation, signed by John E. Fitzgerald, Director	<a href="mailto:jef@jefitzgerald.net">jef@jefitzgerald.net</a>	Yonkers, NY
<b>Individuals</b>		
Aimee Miller	<a href="mailto:Aimee.Miller84@gmail.com">Aimee.Miller84@gmail.com</a>	Mountain Home, ID
Alexandra Croom	<a href="mailto:alexandra@croom.us">alexandra@croom.us</a>	Gales Ferry, CT
Alice Heinzen, Director of the Office for Marriage and Family Life, Diocese of La Crosse	<a href="mailto:aheinzen@diolc.org">aheinzen@diolc.org</a>	La Crosse, WI
Allan Parker, The Justice Foundation	<a href="mailto:parker4justice@gmail.com">parker4justice@gmail.com</a>	
Allison Malkowski	<a href="mailto:allison.suda@gmail.com">allison.suda@gmail.com</a>	Bismarck, ND
Amy Michniak	<a href="mailto:gnamich@hotmail.com">gnamich@hotmail.com</a>	Garland, TX
Andrea Gronsky	<a href="mailto:rongronsky@gmail.com">rongronsky@gmail.com</a>	El Cerrito, CA
Andrea Mack	<a href="mailto:andreaverr@hotmail.com">andreaverr@hotmail.com</a>	Grand Island, NE
Andrea Schanandore	<a href="mailto:taschandy@gmail.com">taschandy@gmail.com</a>	Bismarck, ND
Anita Stevens	<a href="mailto:astevens@polarcomm.com">astevens@polarcomm.com</a>	Cavalier, ND
Ann LaBar Hall, PA-C	<a href="mailto:annlabar@hotmail.com">annlabar@hotmail.com</a>	Atlanta, GA
Ann Makar	<a href="mailto:amakar1@gmail.com">amakar1@gmail.com</a>	Vancouver, WA
Anne Simonnot	<a href="mailto:coueti@hotmail.com">coueti@hotmail.com</a>	Fairfield, CT
Ashley Grunhovd	<a href="mailto:Ashley.grunhovd@fargodiocese.org">Ashley.grunhovd@fargodiocese.org</a>	Fargo, ND
Ashley M. Wiskirchen	<a href="mailto:ashleymwiskirchen@gmail.com">ashleymwiskirchen@gmail.com</a>	Jefferson City, MO
Aurora Verkamp	<a href="mailto:auroraverkamp@gmail.com">auroraverkamp@gmail.com</a>	Brownsburg, IN
Barbara Rose	<a href="mailto:savro@verizon.net">savro@verizon.net</a>	Jenkintown, PA
Betty Lew Coda	<a href="mailto:codafam@aol.com">codafam@aol.com</a>	Honolulu, HI
Bonnie Scheresky	<a href="mailto:agserve@restel.com">agserve@restel.com</a>	Max, ND
Brad Gray	<a href="mailto:brad.gray@fargodiocese.org">brad.gray@fargodiocese.org</a>	Fargo, ND
Brendan O'Connell	<a href="mailto:brendy@verizon.net">brendy@verizon.net</a>	West Roxbury, MA
Brittany Kudrna	<a href="mailto:bnkudrna@gmail.com">bnkudrna@gmail.com</a>	Bismarck, ND
Bruce Bartholomew	<a href="mailto:cabbiemom@gmail.com">cabbiemom@gmail.com</a>	Chugiak, AK
Caitlin Bootsma	<a href="mailto:caitlinbootsma@gmail.com">caitlinbootsma@gmail.com</a>	Richmond, VA
Candy Metoyer	<a href="mailto:cmetoyer@la-archdiocese.org">cmetoyer@la-archdiocese.org</a>	Long Beach, CA
Carmen Devney	<a href="mailto:hoofprince@hotmail.com">hoofprince@hotmail.com</a>	Bismarck, ND
Carrie Huebner	<a href="mailto:carrie@santacasafertilitycare.com">carrie@santacasafertilitycare.com</a>	Canton, GA
Carrie Keating	<a href="mailto:carriekeating@comcast.net">carriekeating@comcast.net</a>	Highlands Ranch, CO
Catherine Schwebach	<a href="mailto:soccerfancsb@gmail.com">soccerfancsb@gmail.com</a>	Lidgerwood, ND
Chelsea Forster	<a href="mailto:chelsealforster@gmail.com">chelsealforster@gmail.com</a>	Bismarck, ND
Chris Reynolds	<a href="mailto:creynolds@ccli.org">creynolds@ccli.org</a>	Cincinnati, OH
Christina Collins	<a href="mailto:ccollins@bismarckdiocese.com">ccollins@bismarckdiocese.com</a>	Bismarck, ND
Christine Bartholomew	<a href="mailto:cabbiemom@gmail.com">cabbiemom@gmail.com</a>	Chugiak, AK



Christopher Bell	<a href="mailto:cxbell@aol.com">cxbell@aol.com</a>	Spring Valley, NY
Cindy Leonard	<a href="mailto:cleonard@dphx.org">cleonard@dphx.org</a>	Phoenix, AZ
Cristofer Pereyra	<a href="mailto:cpereyra@tepeyacleadership.org">cpereyra@tepeyacleadership.org</a>	Phoenix, AZ
Dan Hawrot	<a href="mailto:dhawrot@netzero.net">dhawrot@netzero.net</a>	Steubenville, OH
Daniel West	<a href="mailto:dwest@archatl.com">dwest@archatl.com</a>	Marietta, GA
David Dawson	<a href="mailto:ddawson@arch-no.org">ddawson@arch-no.org</a>	New Orleans, LA
David Prier	<a href="mailto:prierdar@gmail.com">prierdar@gmail.com</a>	Erie, PA
Dawn Becker	<a href="mailto:dawn16marie@gmail.com">dawn16marie@gmail.com</a>	Napoleon, ND
Deanna M. Johnston	<a href="mailto:djohnston@stphilipinstitute.org">djohnston@stphilipinstitute.org</a>	Tyler, TX
Debi Hoppe	<a href="mailto:debihoppe@gmail.com">debihoppe@gmail.com</a>	Irvine, CA
Denise Herding	<a href="mailto:Dherding@msn.com">Dherding@msn.com</a>	Wahpeton, ND
Derek McDonald	<a href="mailto:dmcdonald@rcbm.org">dmcdonald@rcbm.org</a>	Manchester, NH
Donna Dausman	<a href="mailto:dmdausman@comcast.net">dmdausman@comcast.net</a>	Decatur, IL
Dr. Jaime Obst	<a href="mailto:jaimeobst@gmail.com">jaimeobst@gmail.com</a>	Buffalo, NY
Dr. Leroy Huizenga	<a href="mailto:lahuizenga@gmail.com">lahuizenga@gmail.com</a>	Bismarck, ND
Ed Hopfner	<a href="mailto:HopfnerE@SFArch.org">HopfnerE@SFArch.org</a>	El Cerrito, CA
Edward T. Coda	<a href="mailto:edhfs@aol.com">edhfs@aol.com</a>	Honolulu, HI
Elizabeth Elicker	<a href="mailto:elizabeth.elicker@gmail.com">elizabeth.elicker@gmail.com</a>	Phoenixville, PA
Elizabeth Leier	<a href="mailto:elroseleier@gmail.com">elroseleier@gmail.com</a>	Harvey, ND
Elizabeth Parrish	<a href="mailto:Fertilitycareservices@comcast.net">Fertilitycareservices@comcast.net</a>	Perkasie, PA
Emily Cowley	<a href="mailto:ecklump@gmail.com">ecklump@gmail.com</a>	Kansas City, MO
Emily Jacobs	<a href="mailto:jacobs.emilyjean@gmail.com">jacobs.emilyjean@gmail.com</a>	West Fargo, ND
Erika Mayoral	<a href="mailto:jermalog@hotmail.com">jermalog@hotmail.com</a>	La Mesa, CA
Fenar Kashat	<a href="mailto:Fenar1201@gmail.com">Fenar1201@gmail.com</a>	Farmington Hills, MI
Greg Paintner	<a href="mailto:greg.paintner@hotmail.com">greg.paintner@hotmail.com</a>	East Grand Forks, MN
Gretchen S. Lorei	<a href="mailto:gschaberg@gmail.com">gschaberg@gmail.com</a>	Erie, PA
Hannah Barnhorn	<a href="mailto:HBARNHORN@GMAIL.COM">HBARNHORN@GMAIL.COM</a>	Hamilton, OH
Heather Sukut	<a href="mailto:Sukutathome@hotmail.com">Sukutathome@hotmail.com</a>	Rice, MN
Ivan D'Souza	<a href="mailto:idsouza_1967@yahoo.com">idsouza_1967@yahoo.com</a>	Murrysville, PA
J. Budziszewski	<a href="mailto:jbud@undergroundthomist.org">jbud@undergroundthomist.org</a>	Austin, TX
Jacob Loegering	<a href="mailto:jrloegering@gmail.com">jrloegering@gmail.com</a>	LaMoure, ND
James Cherrey	<a href="mailto:jillC@joyfilledmarriagenj.org">jillC@joyfilledmarriagenj.org</a>	Westfield, NJ
Jana Heen	<a href="mailto:jheen9398@gmail.com">jheen9398@gmail.com</a>	Bismarck, ND
Janet Smith, PhD	<a href="mailto:profjanetsmith@gmail.com">profjanetsmith@gmail.com</a>	Detroit, MI
Jassica Heffron	<a href="mailto:jassicadion@yahoo.com">jassicadion@yahoo.com</a>	Colorado Springs, CO
Jayne Stefanic	<a href="mailto:jstefanic@diosav.org">jstefanic@diosav.org</a>	Eden, GA
Jeanne Berdeaux	<a href="mailto:Berdeaux@dioceseofvenice.org">Berdeaux@dioceseofvenice.org</a>	Venice, FL
Jennifer Collins	<a href="mailto:jcollins@rockforddiocese.org">jcollins@rockforddiocese.org</a>	Rockford, IL
Jennifer D'Souza	<a href="mailto:kell.jt@gmail.com">kell.jt@gmail.com</a>	Murrysville, PA
Jeremy Sauer	<a href="mailto:jersaue@gmail.com">jersaue@gmail.com</a>	Max, ND
Jessina Kary	<a href="mailto:ja.designs24@gmail.com">ja.designs24@gmail.com</a>	Dickinson, ND
Jill Cherrey	<a href="mailto:jillC@joyfilledmarriagenj.org">jillC@joyfilledmarriagenj.org</a>	Westfield, NJ
Jillian Stauffer	<a href="mailto:jillianwills@yahoo.com">jillianwills@yahoo.com</a>	Owosso MI
Jo Ann Gerling	<a href="mailto:rjgerling@aol.com">rjgerling@aol.com</a>	New Brunswick, NJ
John S. Grabowski	<a href="mailto:grabowski@cua.edu">grabowski@cua.edu</a>	Washington, DC
Joseph O. DeVet	<a href="mailto:jdevet@archgh.org">jdevet@archgh.org</a>	Houston, TX
Joseph A. Schmidt	<a href="mailto:jschmidt@diosteub.org">jschmidt@diosteub.org</a>	Steubenville, OH

Joshua Schwebach	<a href="mailto:wariorknight817@gmail.com">wariorknight817@gmail.com</a>	Lidgerwood, ND
Julianna Jervis	<a href="mailto:juliannajervis@att.net">juliannajervis@att.net</a>	Reno, NV
Julie Linder	<a href="mailto:julie.linder@dsj.org">julie.linder@dsj.org</a>	San Jose, CA
Karen D. Poehailos, MD	<a href="mailto:cvillenapro@gmail.com">cvillenapro@gmail.com</a>	Charlottesville, VA
Katherine Quigley	<a href="mailto:kaquigley01@gmail.com">kaquigley01@gmail.com</a>	New Haven, IN
Kathleen Robinson	<a href="mailto:kwagsmomof9@yahoo.com">kwagsmomof9@yahoo.com</a>	Lake City, PA
Kathryn Elenchin	<a href="mailto:kelenchin@gmail.com">kelenchin@gmail.com</a>	Coudersport, PA
Katie Monson	<a href="mailto:zilmon23@gmail.com">zilmon23@gmail.com</a>	Williston, ND
Katie Sauer	<a href="mailto:jeresaue@gmail.com">jeresaue@gmail.com</a>	Max, ND
Katrina J. Zeno	<a href="mailto:kzeno@dphx.org">kzeno@dphx.org</a>	Phoenix, AZ
Katrina Welborn	<a href="mailto:katrina.welborn@hshs.org">katrina.welborn@hshs.org</a>	Sturgeon Bay, WI
Kelly Righetti	<a href="mailto:krighetti@srdiocese.org">krighetti@srdiocese.org</a>	Santa Rosa, CA
Kerry Ann Caswell	<a href="mailto:Kac16@cox.net">Kac16@cox.net</a>	Irvine, CA
Kim Elsmore	<a href="mailto:elsmore@dioceseofvenice.org">elsmore@dioceseofvenice.org</a>	Venice, FL
Kristin Detloff	<a href="mailto:detloff.kristin@gmail.com">detloff.kristin@gmail.com</a>	Minneapolis, MN
L. Kwame Fosu, JD/MBA	<a href="mailto:loufosu@gmail.com">loufosu@gmail.com</a>	Washington, DC
Laura Pietkiewicz	<a href="mailto:reading456@gmail.com">reading456@gmail.com</a>	Paxton, IL
Lauren Fuller	<a href="mailto:lfuller@symptopro.org">lfuller@symptopro.org</a>	Portland, OR
Laurie Kraemer	<a href="mailto:motherkraemer@gmail.com">motherkraemer@gmail.com</a>	Grand Forks, ND
Leah Pelto	<a href="mailto:michaelandleah@suddenlink.net">michaelandleah@suddenlink.net</a>	Pineville, LA
Leann Ripplinger	<a href="mailto:fcs.heartland@hotmail.com">fcs.heartland@hotmail.com</a>	Jamestown, ND
Lindsey Marugg	<a href="mailto:Lindsey.marugg@gmail.com">Lindsey.marugg@gmail.com</a>	Buffalo, NY
Lori Hoye	<a href="mailto:life@issues4life.org">life@issues4life.org</a>	Union City, CA
Lucinda Flamiatos	<a href="mailto:lucy@brightdawnnutrition.com">lucy@brightdawnnutrition.com</a>	Lynnwood, WA
Luke Seidling	<a href="mailto:seidling@gmail.com">seidling@gmail.com</a>	Bismarck, ND
Lydia Borja	<a href="mailto:lydia.femm@gmail.com">lydia.femm@gmail.com</a>	Kennesaw, GA
Madeline Wagner	<a href="mailto:Madelinesmiles@gmail.com">Madelinesmiles@gmail.com</a>	Erie, PA
Marguerite Duane	<a href="mailto:MDuaneMD@gmail.com">MDuaneMD@gmail.com</a>	Washington, DC
Maria Prier	<a href="mailto:priermaria@gmail.com">priermaria@gmail.com</a>	Erie, PA
Marisa Merkle	<a href="mailto:Mnsmerkle@gmail.com">Mnsmerkle@gmail.com</a>	Monterey, CA
Marissa Duppong	<a href="mailto:MarissaDuppong@gmail.com">MarissaDuppong@gmail.com</a>	Fargo, ND
Mary Anne Anderson	<a href="mailto:MAanderson@fertilityappreciation.com">MAanderson@fertilityappreciation.com</a>	Orinda, CA
Mary Johnson	<a href="mailto:ryanmaryjohnson@hotmail.com">ryanmaryjohnson@hotmail.com</a>	Moorhead, MN
Mary Pat Van Epps	<a href="mailto:boppie5@att.net">boppie5@att.net</a>	Bartlett, TN
Mary Rice Hasson	<a href="mailto:mhasson@eppc.org">mhasson@eppc.org</a>	Fairfax, VA
Mary Schneider, FNP-BC, Assistant Director, Marquette University College of Nursing, Institute for Natural Family Planning	<a href="mailto:Mary.schneider@marquette.edu">Mary.schneider@marquette.edu</a>	Milwaukee, WI
Mary Wolpiuk	<a href="mailto:stmarsuz@aol.com">stmarsuz@aol.com</a>	
Matthew Wagner	<a href="mailto:Mwagsof7@gmail.com">Mwagsof7@gmail.com</a>	Erie, PA
Melinda Seidling	<a href="mailto:seidling@gmail.com">seidling@gmail.com</a>	Bismarck, ND
Michele Chambers	<a href="mailto:michelebrianc@aol.com">michelebrianc@aol.com</a>	Lincoln, NE

Mikayla Dalton	<a href="mailto:mikayla.mernone@gmail.com">mikayla.mernone@gmail.com</a>	Boston, MA
Molly Marie Daley	<a href="mailto:mollymdaley@gmail.com">mollymdaley@gmail.com</a>	Lutherville, MD
Monica Dvoranchik	<a href="mailto:moe25@hotmail.com">moe25@hotmail.com</a>	Fairview, PA
Monique Kraemer	<a href="mailto:petermoniquekraemer@gmail.com">petermoniquekraemer@gmail.com</a>	Menoken, ND
Mrs. Kari Huizenga	<a href="mailto:leroykari@gmail.com">leroykari@gmail.com</a>	Bismarck, ND
Patrick Flynn	<a href="mailto:pflynn@gw.stcdio.org">pflynn@gw.stcdio.org</a>	Sauk Rapids, MN
Patrick Lee, Director, Center for Bioethics, Franciscan University of Steubenville	<a href="mailto:PLee@franciscan.edu">PLee@franciscan.edu</a>	Steubenville, OH
Patrick Metts	<a href="mailto:pmetts@archatl.com">pmetts@archatl.com</a>	Cumming, GA
Paul M. Lorei	<a href="mailto:gschaberg@gmail.com">gschaberg@gmail.com</a>	Erie, PA
Paul Metoyer	<a href="mailto:paulmetoyer@gmail.com">paulmetoyer@gmail.com</a>	Long Beach, CA
Peter G. Martin	<a href="mailto:pmartin@dowr.org">pmartin@dowr.org</a>	Fountain City, WI
Peter Kraemer	<a href="mailto:petermoniquekraemer@gmail.com">petermoniquekraemer@gmail.com</a>	Menoken, ND
Phil Lawson	<a href="mailto:plawson@vermontcatholic.org">plawson@vermontcatholic.org</a>	South Burlington, VT
Poppy Daniels, MD	<a href="mailto:poppyatdrpoppy@gmail.com">poppyatdrpoppy@gmail.com</a>	Ozark, MO
Rachel Meyer	<a href="mailto:rachelmeyer316@gmail.com">rachelmeyer316@gmail.com</a>	Los Alamos, NM
Richard Fehring	<a href="mailto:richard.fehring@marquette.edu">richard.fehring@marquette.edu</a>	Milwaukee, WI
Ron Gronsky	<a href="mailto:rongronsky@gmail.com">rongronsky@gmail.com</a>	El Cerrito, CA
Ruth Loegering	<a href="mailto:jrloegering@gmail.com">jrloegering@gmail.com</a>	LaMoure, ND
Sandra Volkman	<a href="mailto:sande.volkman@midco.net">sande.volkman@midco.net</a>	Bismarck, ND
Sandy Christiansen	<a href="mailto:SChristiansen.Consultant@care-net.org">SChristiansen.Consultant@care-net.org</a>	Lansdowne, VA
Sandy Driesslein	<a href="mailto:fertilitycarevenice@gmail.com">fertilitycarevenice@gmail.com</a>	Venice, FL
Sarah Sullivan	<a href="mailto:sahara585@yahoo.com">sahara585@yahoo.com</a>	Pearland, TX
Sheila Reineke	<a href="mailto:sreineke@gw.stcdio.org">sreineke@gw.stcdio.org</a>	Clear Lake, MN
Stan Wolpiuk	<a href="mailto:stmarsuz@aol.com">stmarsuz@aol.com</a>	
Stephen Dvoranchik	<a href="mailto:sdvoranchik@hotmail.com">sdvoranchik@hotmail.com</a>	Fairview, PA
Steve Bozza	<a href="mailto:sbozza@cdu.edu">sbozza@cdu.edu</a>	Philadelphia, PA
Steven Mosher	<a href="mailto:steve@pop.org">steve@pop.org</a>	Front Royal, PA
Tara Brooke	<a href="mailto:tbrooke@bismarckdiocese.com">tbrooke@bismarckdiocese.com</a>	Bismarck, ND
Tara Ingebritsen	<a href="mailto:tara_ingebritsen@goodcounselhomes.org">tara_ingebritsen@goodcounselhomes.org</a>	Conshohocken, PA
Terri McCormack	<a href="mailto:tmccormack59@gmail.com">tmccormack59@gmail.com</a>	Rutland, VT
Thomas Lenz, MD	<a href="mailto:tlenzmd@aol.com">tlenzmd@aol.com</a>	
Todd Phillips	<a href="mailto:tphillips@gbdioc.org">tphillips@gbdioc.org</a>	Appleton, WI
Tom O'Neill	<a href="mailto:toneill@hbgdiocese.org">toneill@hbgdiocese.org</a>	Mechanicsburg, PA
Verónica F. Arias	<a href="mailto:ariasv@archspm.org">ariasv@archspm.org</a>	Saint Paul, MN
Walter B. Hoyer, II	<a href="mailto:life@issues4life.org">life@issues4life.org</a>	Union City, CA
Xhonane Olivas	<a href="mailto:xolivas@yahoo.com">xolivas@yahoo.com</a>	Clarion, PA